U. S. DEPARTMENT OF AGRICULTURE.

FARMERS' BULLETIN No. 104.

Has been rev.
--see rev.ed.
binders at
end of file.

NOTES ON FROST.

BY

E. B. GARRIOTT,
Professor of Meteorology, Weather Bureau.

WASHINGTON:
GOVERNMENT PRINTING OFFICE.
1899.

LETTER OF TRANSMITTAL.

U. S. DEPARTMENT OF AGRICULTURE,
WEATHER BUREAU,
Washington, D. C., June 15, 1899.

DEAR SIR: In compliance with directions contained in your letter of March 29, 1899, I have the honor to transmit herewith a paper on the subject of frost, prepared by Prof. E. B. Garriott, Weather Bureau, and to recommend its publication as a Farmers' Bulletin.

In an article published in 1896 in Weather Bureau Bulletin No. 86, Prof. W. H. Hammon furnished valuable information regarding methods of protection from frost which have been tested and employed in orchards, and more especially in the citrus fruit region of California.

In the autumn of 1896 the California State Board of Horticulture published a more extensive bulletin, which embodied the experiments and suggestions contained in Weather Bureau Bulletin No. 86, and treated, in an exhaustive manner, the problem of protecting California orchards from frost.

In Weather Bureau Bulletin No. 23, 1899, Professor Hammon presents a revised edition of the bulletin formerly prepared by him and submits modifications of many of his former statements which are made necessary by more recent experiments.

The methods advocated in the bulletin referred to were mainly the result of experiments conducted in California orchards, while a bulletin on frost, applicable, as far as possible, to all localities rather than to restricted districts, would doubtless be of great value to farmers, fruit growers, and gardeners; hence your direction that the Weather Bureau furnish for publication as a Farmers' Bulletin a paper on frost which shall in a popular way define frost and the conditions which favor its formation, and indicate methods of protection which have been found practicable by actual experiments. This suggestion the author has endeavored to carry out in this bulletin.

As the California experiments quoted in Bulletin No. 23 were perhaps the most thorough ever made in this country, a description of the methods employed and the results obtained are reprinted in brief in this paper, together with results of similar experiments made in the citrus fruit regions of the Gulf States, where varying climatic conditions have been found to produce somewhat different results.

Following this summary the subject of frost protection as it relates to the various other agricultural products of the country is considered, and some facts regarding freezes, which are destructive alike to tender vegetation and to plants of hardier growth, are presented.

Your wishes, as communicated in the letter referred to, that the subject be treated in a popular way and divested of all technicalities, and made applicable to all localities rather than to restricted districts, have been complied with so far as the nature of the subject will permit.

I am, very respectfully,

WILLIS L. MOORE,
Chief of Weather Bureau.

Hon. JAMES WILSON,
Secretary of Agriculture.

CONTENTS.

	Page.
How frost is formed	5
Seasons of frost	6
When to expect frost	7
Protection from frost	10
Devices for preventing rapid radiation of heat	11
Glass screens	11
Cloth screens	11
Lath screens	11
Other methods	11
Smudge fires	12
Devices for adding moisture to the air	12
Fires of damp straw and stable manure	13
Sacks of manure	13
Bales of wet straw	13
Prunings	13
Portable smudge fires	13
Devices for charging the air with moisture	14
General observations and practical results	17
Work in Texas	17
Work in Louisiana and Florida	19
Frost seasons for several crops	21
Conclusions	22

NOTES ON FROST.

HOW FROST IS FORMED.

The atmosphere of the earth always contains more or less moisture in an invisible form. When at a considerable elevation above the earth this moisture, or aqueous vapor, is condensed, clouds are formed; when the process of condensation is more active and the temperature of the air is above freezing, rain falls; and when the temperature of the air is below freezing, snow is produced. When the moisture of the air in immediate contact with the earth is condensed at temperatures above freezing, dew is formed; when at temperatures below freezing, frost is deposited. Frost is, therefore, the moisture of the air condensed at freezing temperatures upon plants and other objects near the surface of the earth.

In the process of frost formation the temperature of the air a few feet above the earth is commonly several degrees above freezing. The surfaces upon which frost is deposited must, however, possess freezing temperatures. The manner in which frost is deposited on plants and other objects is very similar to that observed when the air moisture of a room is frozen and deposited upon window glass, the temperature of which has been reduced to freezing by the out-of-doors cold. In the case of the frost on the window glass the process is one that can readily be understood. Some explanation, however, is required of the formation of frost, which requires freezing temperature, when at times a temperature *above freezing* is registered a few feet above the surfaces upon which frost appears.

There are several processes by which the temperature of plants may be reduced below the temperature of the air which surrounds them. The most important of these processes is radiation, by means of which heat escapes from objects and passes into the surrounding air.

In the frost-forming process heat from the sun which is absorbed by the earth and by plants during the day is lost by radiation at night. During the day the earth both absorbs and reflects the heat received from the rays of the sun, and the lower stratum of air is warmed by this reflected heat. During the night when no direct heat is received from the sun's rays the lower air stratum receives no reflected heat, and, at the same time, heat which has been absorbed by the earth is radiated, or, in other words, it rises through the overlying air, though not

necessarily and entirely as sensible heat; that is, heat perceptible to the senses. In the frost-forming process there is another very important factor in the production of cold; that is, the evaporation of moisture from the earth and from plants. The nature and composition of frost require that the air in immediate contact with the surfaces upon which frost forms shall contain moisture. This is moisture that has been stored in the earth in visible form, i. e., in the form of water, and which has escaped from the earth, not as visible moisture, but by the process of evaporation and in the form of aqueous vapor. Some part of the heat stored by the earth during the day is, therefore, lost in evaporating the moisture contained in plants and in the earth. The heat utilized in this process is retained in the air, not as sensible, but as latent heat. Heat is rendered latent, in the same manner, when it is used to boil away water, and it again becomes sensible or liberated heat when the water vapor produced by boiling, which is a rapid process of evaporation, is again condensed into water. It appears, therefore, that while the temperature of surfaces upon which frost forms, and of the air in immediate contact with them, is lowered by the evaporation of moisture from the surfaces, the influence of the process does not extend to air a few feet above the ground.

Another method by which plants lose their heat is by convection, by means of which they are chilled by contact with colder air. This process, while important in the presence of freezing-air temperatures, can scarcely be considered a factor in the formation of frost proper, which is usually accomplished when temperature observations show the air to be above freezing.

Another element in the formation of frost is found in the fact that air, like fluids, arranges itself according to its density or weight. Thus the air in immediate contact with the earth becomes heavier as its temperature is lowered by the radiation of heat and settles into depressions or valleys and over lowlands, causing, at times, frost in low-lying districts, while neighboring higher grounds escape the visitations.

SEASONS OF FROST.

Generally speaking, agricultural products in some part of the United States are menaced by frost from the time nature begins to extend her mantle of green from the citrus regions of the South toward the grain fields of the North until the verdured area contracts southward before the first winds of winter; from the hour that seeds germinate and buds blossom until crops are gathered; from early spring until late fall. As a rule, the critical periods are during the early growth of plants and when the fruit is in the bloom, and again when crops are maturing. In the first instance vegetation is threatened by the frosts of early spring, and in the latter by the frosts of autumn.

Considering the several agricultural sections in greater detail we find that damaging frost is likely to occur in the north half of the Florida

peninsula and in the region immediately bordering the coast of the Gulf of Mexico, and in the Pacific Coast States from the latter part of October until the early part of April, and in the North Pacific Coast States as early as the middle of October and as late as the last week of April. From December until early in March the frost limit may, at long intervals, be extended southward well over central and even over southern Florida. The period of damaging frosts in the interior of the South Atlantic and Gulf States extends from November to April. From October to April the region in which agricultural products are subject to damage by frost is extended to the southern lake region, over the Upper Mississippi and Lower Missouri valleys, Kansas, and Nebraska, and from September to May frost visitations are usually confined to the extreme Upper Mississippi, Upper Missouri, and Red River of the North valleys, and the Rocky Mountain and plateau regions from central New Mexico and the Texas Panhandle northward. During the period, June to August, inclusive, frosts are very infrequent, and when they do occur are confined, practically, to mountain districts and to the northern tier of States.

WHEN TO EXPECT FROST.

As the conditions which produce damaging frost are subject to modifications which are as numerous as the kinds of crops raised, and as varied as local topography, local climate, and local soil conditions in the various sections of the United States, the writer in attempting to treat briefly a subject which is so diversified in its aspects is confronted with many difficulties. Suffice it to say that with other atmospheric conditions favorable for its occurrence frost may, as a rule, be expected when temperature, as reported by stations of the Weather Bureau, falls to a point 8° to 10° above the freezing point. As previously stated, the surfaces upon which frost is deposited must possess freezing temperature, although the temperature of the air a few feet above the surfaces may be several degrees above freezing; and it is the temperature of the air, in some instances many feet above the ground, that is given by the Weather Bureau observations. Another atmospheric condition favorable for the occurrence of frost is a clear, cloudless, and comparatively calm night. The presence of clouds retards radiation or loss of heat from plants; the clouds act as a screen in preventing the heat collected from the sun's rays during the day from escaping into the upper air. When clouds are not present and a withdrawal of the sun's rays causes a rapid cooling of the air at moderate elevations, the warmer air which collects near the surface of the earth during the day rises, and the cooler upper air, owing to its greater density or weight, settles to the earth. It will be noted that clouds not only prevent the escape of the warm air in immediate contact with the earth, but also blanket, as it were, the upper part of the lower air stratum. Calm or comparatively still air is a condition which favors

the formation of frost. On windy nights the air is disturbed and is not permitted to arrange itself in layers according to its density, with the densest and coldest air near the surface of the earth; it is kept mixed up by the wind.

A very interesting fact, which illustrates the manner in which cold air settles by its greater weight to the surface of the earth and allows the warmer air to assume higher levels, causing frost in low grounds and granting to the higher grounds a comparative immunity from frost, is set forth in the following article, by Prof. Cleveland Abbe, in the Monthly Weather Review for December, 1893, entitled Thermal Belts, Frostless Belts, or Verdant Zones:

These are local names given to certain regions on mountain sides within which nocturnal frost rarely or never occurs in the springtime, although freezing temperatures occur in the winter time, consequently tender vegetation flourishes with remarkable vigor in these regions. The following are the only references to thermal belts in the United States at present known to the editor, but as such regions are specially important to the horticulturist and agriculturist it is hoped that the correspondents of the Weather Bureau will bring these thermal belts to notice wherever they occur in order that their meteorological peculiarities may be better understood.

In the Agricultural Report of the Patent Office for 1861 Mr. Silas McDowell, of Franklin, Macon County, N. C., describes the verdant zone in that county in the valley of the Little Tennessee River. It occupies the region between 300 and 700 feet above the valley of the river, which latter is about 2,000 feet above sea level. On tracing this zone up among the smaller tributaries of the Tennessee River he found that in the higher valleys, where the bottom land is about 3,900 feet above sea level, the verdant zone lies between 4,000 and 4,100 feet. Within this zone frost never injures the vegetation, and the most tender grapes never fail to produce abundant crops.

Prof. J. W. Chickering, jr., in the bulletin of the Philosophical Society of Washington, March, 1883, and in the American Meteorological Journal, vol. 1, describes the following thermal belt: "In Polk County, N. C., along the eastern slope of the Tryon Mountain range, in latitude north 35, the thermal belt begins at the base of the mountain, at an elevation of 1,200 feet above the sea, and extends up 2,200 feet, being most perfect at about 1,500 feet. It is about 8 miles long, and is distinguished by magnificent flora, such as would be characteristic of a point 3 degrees south of the actual latitude."

Prof. John Leconte, of Berkley, Cal., in Science, vol. 1, p. 278, states that at Flat Rock, near Hendersonville, Henderson County, N. C., on the flank of the mountain spur adjacent to the valleys of the Blue Ridge, he also observed a frostless zone. The valley is about 2,200 feet above sea level, and the thermal belt is 200 to 300 feet above the valley.

Mr. J. W. Pike, of Vineland, N. J., states that among the mountains of California he has discovered that during the night the cold is much greater in the valleys than on the terraces several hundred feet above, due to the settling of the cold air, so that a thermal belt is formed at that height separating the frosty valleys from the colder highlands.

In the Tennessee Journal of Meteorology for January, 1894, published by the State weather service, the author describes a thermal belt between Los Angeles and the Pacific coast. It traverses the foothills of the Cahuenga range, and has an elevation of between 200 and 400 feet and a breadth of about 3 miles. It occupies the midway region of the range.

In the American Meteorological Journal, vol. 1, Mr. S. Alexander describes a thermal belt in which the peach tree flourishes in the southeastern portion of Michigan. He shows that the cold island discovered by Winchell in that region is really the bottom of a topographical depression into which the cold air settles. It is a long valley surrounded by a belt of elevated country from 50 to 600 feet above lakes Michigan and Huron. The valley and the isotherms trend northeast and southwest from Huron County through Sanilac, Lapeer, Oakland, Livingston, and Washtenaw to Hillsdale counties. The highlands of this region are all much freer from frost than the lowlands, and all much more favorable for early vegetation. He does not state that any point is high enough to be above the thermal belt, but that, in general, two equal parallel thermal belts inclose the cold island between them.

It is generally conceded that these thermal belts depend both upon the drainage of cold air downward into the lower valleys and the freedom of radiation from the surface of the ground to the clear sky overhead. During a still night, when frosts occur, the surface of the hillside cools by radiation, and hence cools the air in contact with it; the latter flows downward as long as its cooling by radiation and conduction exceeds its warming by compression. Inasmuch as its cooling depends on contact with a still colder soil or plant, it soon accumulates in the lowlands as a layer of cold air, which grows thicker during the night by the steady addition of the thin layer of descending air in contact with the ground on the hillsides. The warmer air, which has not yet had an opportunity to cool by contact with the ground, floats on top of the cold mass; it spreads out toward the hills, and is continuously furnishing its heat to the adjacent hillsides as fast as it comes in contact with them before it also cools and descends. The formation of the thermal belt seems to depend largely upon this gentle circulation during the nighttime. The lower limit of the belt is defined by the depth of the accumulation of cold air in the confined valley and rises higher in proportion as the night is clearer and longer, and also in proportion as the valley is more or less perfectly inclosed. The upper limit of the thermal belt may depend upon the strength of the wind and the general temperature of the air. But if there be no wind, then it depends equally on the freedom of radiation to the clear sky and on the above-described circulation of air.

The facts quoted in this article show that frost may be expected on low grounds at times when higher grounds escape the visitations, and the lesson they teach is that early and tender crops should, so far as may be practicable, be confined to crests, hillsides, and mountain sides, and later and hardier crops to the lowlands and valleys.

Local climate, as it is influenced and regulated by the proximity of bodies of water, must be given great weight in calculations regarding frost. Frost is less likely to occur in localities swept by moisture-laden air which has crossed a considerable body of water. This is more especially the case in the fall of the year, when the temperature of bodies of water is reduced very slowly; and in the South, where the water temperature continues relatively high throughout the year, the influence of the water is especially marked. During the colder months air is not only warmed in crossing considerable bodies of water, but it also absorbs moisture, which, although invisible in the form of water vapor, has the effect of retarding the radiation of heat from the earth. In fact, so pronounced is the influence of water and water vapor that localities which with reference to exposure to west and northwest winds are, as it were, in the lee of large bodies of water are comparatively free from frost visitations. Along parts of the east Florida

coast which are protected on the west by the Indian River, frost is said to be unknown, and throughout the central and northern sections of the United States, localities which are protected on the west and northwest by bodies of water of considerable size are not only favored, by reason of their position, with conditions which do not promote the formation of frost, but also receive the benefits derived from heavier falls of snow.

Local soil conditions constitute, to a certain extent, a factor in the formation of frost. As a rapid loss of heat is promoted by an active evaporation of moisture, it is evident that, with other conditions equal, frost is more likely to occur on damp than on dry ground, provided, however, that the ground is not too moist, for in the latter event the amount of moisture evaporated and added to the air would have a tendency to retard the radiation of heat from the earth. And herein lies the distinction, so far as frost formation is concerned, between moist air and moist soil. An excess of moisture in the air, in preventing, to a degree, the radiation of heat, is unfavorable to the formation of frost. As frost is the moisture of the air in immediate contact with the earth condensed at freezing temperature, it follows that the earth, from which the moisture of the air is drawn, must contain more or less water, and it is evident that damaging frost will occur with a limited rather than with a large quantity of moisture arising from the earth and a moderately moist and comparatively still air. Both of these conditions usually obtain following, but not too closely following, the rains of spring and autumn. The character of the soil as regards its capacity for retaining moisture must also be taken into account, more especially during periods of comparatively dry weather. Moist soil, and the vegetation which springs from it, is chilled to a lower temperature by the evaporation of moisture, and is therefore more subject to visitations of frost than soil which is dry, or into which water penetrates deeply, leaving the surface dry, or soil which sheds the rain and does not absorb and hold water. It is important to note, however, that very moist soil, or soil which contains a large amount of surface water, is, owing to the quantity of moisture yielded to the air, not so subject to the heavy and damaging frosts which visit plants on moderately damp ground or ground that has a small capacity for moisture.

PROTECTION FROM FROST.

A discussion of methods of protection from frost calls for a consideration in great detail, not only of the several approved processes of protection and the extent to which they can be applied in connection with the various crops, but also the value of these processes under the varying conditions of local topography, local climate, and local soil conditions.

It is evident, in the light of the foregoing statements, that fruit and vegetable growers in rolling and hilly country, or when located in the lee, as regards west to northwest winds, of bodies of water, can, to some

extent, so place their earlier and tenderer fruits and crops that they will, in the first instance, avoid the lowlands and valleys, and in the second take advantage of the more moderate and moist air that crosses water surfaces. For further protection the grower must depend upon artificial appliances. These appliances are designed to produce the following effects or results: To prevent a rapid radiation of heat from the earth; to charge the air with moisture; to warm the air; to create artificial drafts, whereby the air is mixed and the cold air is not allowed to settle to the surface of the earth; or to actually cover or roof in plants.

DEVICES FOR PREVENTING RAPID RADIATION OF HEAT.

Devices designed to prevent a rapid radiation of heat from the earth include screens which can be drawn over plants, vineyards, and groves; loose substances with which low plants may be covered; and smudge fires built to the windward of areas for which protection is desired. These devices, which of a necessity are operative for very limited areas, are described by Professor Hammon in Bulletin No. 23, Weather Bureau, as follows:

Glass screens.—In greenhouses and hotbeds advantage is taken of the peculiar property of glass which allows the heat rays of the sun to pass through it, and is almost impervious to the dark *heat* rays from the earth and plants. This is one of the most perfect screens possible, since it not only prevents the loss of heat by radiation, but receives and retains the heat from the sun. The expense precludes its adoption except for the protection of valuable plants and flowers.

Screens of other solid materials have been quite extensively used in protecting vineyards and citrus groves where intense cultivation is practiced and where the location of the groves near an excellent market admits of profit even with expensive methods of cultivation.

Cloth screens.—In Italy and portions of France screens made of muslin strung on wires stretched on poles above the tops of trees or vines have been used extensively. These screens are drawn on nights when frosts are probable, and are pushed back during the day. When the season has advanced so far as to preclude further danger, they are taken down and stored. Of course such a plan could be operated only on a very limited scale, and would then be expensive. This plan has been recently successfully tried in the orange groves of southern California.

Lath screens.—During the past few years screens made of laths fastened to ordinary telephone wire (the spaces between them being about the width of the laths) have been extensively used in Florida. These are spread over a frame erected above the trees or plants. The screens serve not only as a fair protection from frost, but also as a shade from the hot sun. When no longer needed, they can be rolled up and stored away for preservation. At first thought it would seem improbable that a screen covering only half the space (the spaces being as wide as the laths) would afford much protection; but when it is considered that laths have considerable thickness it is plain that while only one-half the vertical rays are screened those inclined between the vertical and horizontal are partially intercepted by the edges as well as the faces of the laths. As a matter of fact, about three-fourths of the sky is screened by this means. By placing the laths in north and south directions the direct rays of the morning sun are completely cut off from the orchard, which admits of the temperature rising slowly. This reduces the liability of injury to plants.

Other methods.—Strawberries and other low plants are frequently protected by covering them with straw or other loose substances.

Frequently young potato plants are saved by plowing a furrow alongside and allowing the dirt to bury them.

Cranberry growers in the marshes of Wisconsin flood the marshes when frost is expected. In this case the protection is probably due, for the most part, to the high specific heat of the water, as only portions of this land are submerged.

Smudge fires.—Since on cloudy nights radiation is so reduced as to prevent the formation of frost, many have thought that an artificial obscuration of the sky by means of dense smoke would be an excellent means of protection. The efforts of this character which have been made have resulted in decidedly varying success. In the wheat fields of the Dakotas excellent protection was obtained in limited areas, while the experience of orchardists in Florida and southern California has not shown such uniform success.

Since it was supposed that the protection resulted from the obscuration of the sky by means of smoke, the best protection was expected from the use of that fuel which would produce the greatest smoke.

In the Dakotas the best and most convenient material at hand was the straw of the previous year's crop, which had been left in the fields all winter and through the rainy spring, until it was quite thoroughly soaked with water.

In southern California and Florida straw was scarce, and where it could be obtained it was much drier than that used in the Dakotas. Consequently tar, crude petroleum, and other similar smudge materials were substituted; but the results have not been, as a rule, satisfactory, although the smoke was equally dense. However, quite successful results were obtained by Mr. Buck, Mr. La Rue, and others in the Vacaville and Sonoma sections by burning damp stable manure in sacks scattered throughout the orchard.

Devices for preventing a rapid radiation at night of the heat absorbed by the earth during the day can be utilized only for comparatively small areas. The screen devices, also, by reason of their cost, are available only for the smaller orchards and vineyards whose products command early market prices; the same may be said of coverings of straw and other materials for strawberries and other low plants. Smudge fires can be used to advantage for orchards, vineyards, and ground plants, and even for the smaller grain fields, and would be particularly efficacious in protecting crops and plants in low or bottom lands over which, on still nights, the smoke from smudge fires would settle. There is no device at present known that will, by preventing a rapid radiation of heat, afford any material protection to the great grain fields of the West and Northwest.

DEVICES FOR ADDING MOISTURE TO THE AIR.

Smudge fires made of damp material have been more successfully used for the purpose of affording protection from frost than those made of tar, crude petroleum, and other dry materials, for the reason that in addition to creating a dense smoke they add to the atmosphere a considerable amount of evaporated water, which, though invisible, serves to retard the radiation of heat. When any considerable amount of water is thus converted into vapor and is distributed in the air which covers the area to be protected, a portion of it is likely to be condensed by the surrounding cooler air and appear in the form of mist, which

acts as an agent to prevent the escape of heat from the earth; and the act of condensation sets free in the air some of the heat that has been expended in the process of evaporating the water contained in the smudge-fire materials. Thus the process is well ordered throughout. In the first place, an undue quantity of heat, whereby undesirable drafts would be created, is not communicated to the air by the fires, and the final result is that in reliberating, by condensation, the water in the air the particles of moisture remain suspended in the air, and some of the heat of the fires is returned to the air in a form which, to a certain degree, warms it without causing currents which would be calculated to disturb the smoke cloud, or pall, and the moist air which overlie the ground and upon whose presence the safety of the plants from frost depends.

As smudge fires made from damp materials are one of the recognized devices which have been found to be more or less effective in affording protection from frost, the following description of methods which have been adopted for utilizing them in orchards is quoted from Bulletin No. 23, already referred to:

Fires of damp straw and stable manure.—Have the fuel, in small piles, distributed throughout the orchard in advance; the more numerous the piles the better. With the same amount of fuel the best protection is obtained from small and frequent fires, since with small fires the upward draft is reduced to a minimum, and the more frequent the fires the more uniform will be the distribution of heat.

Sacks of manure.—A decidedly preferable method is to pack damp stable manure in common grain or burlap sacks, by which it can be conveniently handled. They should be distributed through the orchards in rows of about 100 feet apart and about 50 feet between sacks in each row. When it is found necessary to protect, a small amount of coal oil is poured upon each sack and ignited. It is usually unnecessary to fire more than every second or fourth sack, the remainder being left for later occasions. These sacks will burn with a smoldering fire for several hours.

The amount of heat which is set free by burning one sack of manure weighing about 50 pounds and condensing the water vapor near the earth would be sufficient to raise the temperature 20° in a space 75 feet square and 25 feet deep. If one-fourth of this heat remained within this region needing protection, which seems to be a reasonable estimate, ample protection would be obtained for almost any ordinary conditions.

Bales of wet straw.—Mr. T. A. Morrison, of Riverside, Cal., suggested the use of a similar plan, in which bales of wet straw were substituted for manure. This plan has been tried with fair success. One hundred pound bales were cut in four pieces, a tie wire being left about each piece, and if properly dampened will burn with but little care, causing a small smoldering fire.

Prunings.—The prunings of the trees, which are usually removed shortly before the period when frosts are likely to do their greatest injury, are excellent smudge material, and should always be preserved for this use. They should be piled in open spaces throughout the orchard or vineyard and burned at times when protection may be needed. The best results will be obtained from as small fires as will result in burning the prunings.

Portable smudge fires.—A number of excellent devices have been tried, in which the fires were built upon some vehicle by which they could be moved about the orchard. The advantages of this plan are several.

First. The fire can be moved to the section where most needed, which is generally along the windward side of the orchard.

Second. The loss of heat by an upward draft is almost entirely prevented, since the fire does not remain in one position long enough to establish such a draft. On this account much larger and, consequently, fewer fires, with equal efficiency, are possible.

Third. There is a much more uniform distribution of heat and smoke throughout the orchard.

As smudge fires prepared and applied in the manner above indicated are inexpensive, to a degree effective, and easy of operation, they constitute the method which for practical use and utility seems best adapted for general use in orchards, vineyards, gardens, truck farms, and small fields, as a protection from frost. Among the many devices that have been employed for holding and carrying these fires the following has been used with especially satisfactory results by the Fleming Fruit Company, of Visalia, Cal., and will doubtless prove equally effective in other sections:

Wire frames (chicken-yard fencing) are built on low truck wagons and stretched from four wagon stakes and heaped over with wet manure. Dirt is then thrown on the wagon beds to protect them, and pots of burning tar are set underneath the straw roof. A barrel of water on the wagon is used to keep the straw wet. The wagons are driven about and do the best work, as they can go wherever the most needed. The smoke and vapor are carried to the rear as the wagon moves, and, being carried at once out of the rising heat, fall close to the ground in a long, white trail. At daylight 400 acres of orchard are covered with a white fog, extending from the ground to about 20 feet high. Similar fires are used as stationary smudges, the wire netting being stretched between four stakes driven in the ground.

Crude petroleum, supplied to a burner from a tank on a wagon or truck, has been used with success by Mr. R. H. Howard, of Riverside, Cal. Over the burner is placed a large wire basket containing mineral wool, which is kept saturated with water supplied by a tank, also placed upon the wagon. Wet straw and manure can also be used. The apparatus can be driven where most needed. The burning coal oil gives off a dense smoke, while the wet mineral wool furnishes a vast surface from which evaporation can take place, and at the same time the material will not be consumed.

DEVICES FOR CHARGING THE AIR WITH MOISTURE.

Aside from the smudge fires, Bulletin No. 23 quotes devices which have been adopted to charge the air directly with moisture, some of which are constructed on a plan too elaborate and expensive for general use.

The first operation involved is the evaporation of water in large quantities, either by means of evaporating pans or by the use of boilers with connecting pipes whereby steam can be generated, carried, and thrown into the air in different parts of a field or orchard.

Spraying and sprinkling plants in times of threatened frost has also been attended with some degree of success. At the Everest ranch,

at Riverside, Cal., sprinklers are placed at the top of 50-foot masts, which fill the air with a very fine spray. In nearly every instance the protection against frost by this method has been sufficient.

Sprinkling gardens before sunrise on frosty mornings has proved of decided value.

Secondly, irrigation, which of necessity can be used only to a very limited extent and in certain favored localities, has been found to afford very complete protection from frost in fields, orchards, and vineyards which are equipped with irrigation ditches, and in the cranberry districts of Wisconsin perfect protection from damage by frost is secured by flooding the cranberry marshes the day before heavy frost occurs in that section.

In no case should water be allowed to stand in orchards, vineyards, or gardens during the early growing period while danger from frost is possible, for the reason that water unduly promotes the growth of young plants and thereby renders them more susceptible to damage by frost and cold than are the plants whose growth is retarded.

The third operation involved in the California experiments was a direct heating of the air by means of fires. It was found that large fires caused a strong upward draft of air, which had the effect of drawing in cooler air from the sides, and that well distributed small fires were attended by very satisfactory results.

The following final report of the Riverside Horticultural Club covers the results of experiments made during the winter of 1897–98:

With the assistance of fifteen or twenty citizens interested in the study of the points involved, a most complete test has been made of the many different methods employed to prevent frost damage. With such a force of competent and impartial observers it was possible to secure data of much value in forming an estimate of the efficiency of the various plans made use of. Careful comparison was made between those orchards where no work was done, and where no direct effect of the fire was probable, and those where the different methods were being tried. As indicated by our partial report at the last meeting of the club, these tests were in some particulars eminently satisfactory as showing the way to definite conclusions.

The exceptionally long period of cold following gave additional opportunity to verify the first conclusions reached, and subsequent investigations made by ourselves, as well as by other citizens who have awakened to the possibility of protecting their property, strengthened and confirmed the opinion formed as the result of the tests already partially reported upon.

Some theories are proved to have little practical value, and members of your committee have modified their views somewhat in consequence. No preconceived notions have been allowed to stand in the way of a thoroughly practical study of the facts as they exist, to the end that the growers may not, for the lack of definite knowledge as to the directions their efforts should take, neglect reasonable precautions hereafter to insure the safety of their crops. These, however, are our conclusions:

First. There is no doubt whatever that the temperature of our orchards may be materially raised by the use of dry heat.

Second. The radiation of the earth's heat can be considerably lessened by moist smudges, when these are started early enough and are properly managed.

Third. The possibility of raising the dew point on one of the dry, cold nights peculiar to our climate sufficiently to prevent damage, by means of steam-producing apparatus, seems impracticable.

Fourth. Fruit and trees can undoubtedly be saved, even in the coldest sections, by covering them with cloth or matting, but the expense involved makes this method impossible on the part of the ordinary grower.

Fifth. It is found that the temperature in an old seedling grove, or where tall wind-breaks afford to smaller fruit trees a like protection, the temperature is almost invariably 1° to 2° higher than in exposed orchards in the immediate neighborhood. This fact seems to thoroughly upset the theory, strongly held by many intelligent growers, that the tall, well located wind-break is a disadvantage, the contrary seeming to be the truth.

Sixth. It is found that the temperature 20 feet above the ground is 1° to 2° higher than at the surface, and that, as a rule, when the cold is severe enough to injure the ripest fruit 50 feet from the ground there is almost invariably a temperature above the freezing point of water.

Professor Zumbro, who has given especial attention to this matter, finds that at the height of 50 feet the temperature is from 5° to 10° higher than at the surface when the air is not in motion. When there is any considerable breeze it varies but little.

Seventh. Our conclusion is that, all things considered, the coal baskets, sufficiently numerous, will prove the most satisfactory and effective means of warming the orchards yet made use of. It is true the oil pots make a far hotter fire, and are neither expensive nor difficult to manage; but the deposit of lamp black upon tree and fruit resulting from their use condemns this system for general use.

As to the value of smudging the members of your committee are not so well agreed. Because of less sharply defined results, we find it more difficult to come to definite, at least uniform, conclusions. But, under certain conditions, we are convinced that, properly used, it may be made a valuable means of protection. We think this especially true in localities where the temperature falls only a little below the danger point, and where there are considerable solid areas of young orchards exposed. Here it will work well if the protection is made general. But where the danger is considerable we think it wise to be prepared to use dry heat, even where in connection with the smudge. The benefit from smudging is probably as much from its protecting fruit and trees from the sudden rays of the morning sun after a freezing night as from modifying temperature during the time of danger.

Experience demonstrates that flooding or running water in connection with dry heat or smudging is a valuable adjunct. One of the committee who has been testing this matter carefully for three years is disposed to think that the direct benefit from running water is overestimated by a majority of growers. Its value in putting orchards in condition to withstand quite severe weather safely is questioned, but the committee are inclined to think that entire dependence upon this method will occasionally result in serious loss to those who trust to this means alone, especially when used in young orchards.

As to the number of baskets needed when coal is used, we find that the most decided and satisfactory results have been gained where from twenty to fifty coal fires have been used to each acre. If intelligently and energetically used this plan will never fail, except when the mercury drops below 24° for a long while, and even then it is believed the larger portion of the crop may be saved if anything like a general use of such fires be secured. The smaller number of fires named has, in numerous instances, and even when a man was working alone, secured a rise of from 3° to 5°, and saved a crop. Can it be doubted that 50 fires per acre, used in every orchard, would save both trees and crop on the coolest night ever known in California's history?

To equip an orchard with 50 baskets to the acre means an outlay of only a little over $5. The fuel to run them one night costs from $2.50 to $3. If a crop of navels upon it is worth $400 it will pay well to spend in labor and fuel $4 per night, or 1 per cent on the value of the crop to insure its safety. In the orange region of southern California it is not unusual to have more than two or three nights in a season when the fruit is in danger. But even if, as in the present season, the period of cold is more extended will it not pay to expend at least as much as one pays for his irrigating water to insure the safe maturing of a crop it has cost him a year's labor and heavy expense to produce? The conclusion is obvious that we have only to provide for the insurance of this sort of property exactly as we would in the case of that liable to destruction by fire, to be enabled to follow the business of orange and lemon growing with the certainty of having perfect fruit to market at the season's end.

While the practicability of protecting our orchards from frost seems established, the problem of the most economical and scientific means of accomplishing this is probably yet to be solved. However well the wire baskets may serve us now, there will doubtless be improved methods for burning coal, and even other material may be found that will serve the purpose better; and while wet straw seems to be at present the most available for smudges, doubtless, when the need is made known, chemists will find some vapor-producing material more compact, efficient, and economical. Hence we recommend that the club appoint a permanent committee to continue these investigations.

GENERAL OBSERVATIONS AND PRACTICAL RESULTS.

WORK IN TEXAS.

In a paper read before the annual meeting of the Texas Farmers' Congress and Texas State Horticultural Society in joint session at College Station, Tex., July 26, 1899, Mr. Joseph L. Cline, observer, Weather Bureau, submitted the following remarks on "The use of frost and temperature warnings in protecting fruits and truck gardens:"

It has been proved that a good way for protecting berry crops is to cover with hay. Berries are improved by mulching. Prairie hay used for this purpose can be cut and placed in the fields, ready for covering the plants when frost is indicated, at a cost of about one dollar a ton. Ten tons per acre are considered sufficient for protection against any freeze. The cost of labor for covering the plants would be about two dollars an acre additional. As the hay is required for mulching, the cost is considerably lessened when the hay is utilized for that purpope. Berry crops which were covered prior to the freeze of last February yielded four times as much as unprotected plants, and unprotected plants were generally killed. Cabbage, potatoes, and other vegetables can be protected from frost by hay, and the increased yield will more than pay for the expense involved. Rice growers along the east Texas coast can raise vegetables throughout the winter on their rice plantations by protecting them from frost. The plantations can easily be flooded during freezing weather, which is always of short duration, and the water would protect and do very little or no damage to the vegetables, which are remunerative during the winter season.

* * * * * * *

The method of burning stakes soaked in petroleum or kerosene oil throughout the fields can be applied not only to the protection of maize, but fruit orchards as well. By means of burning stakes fields will be enshrouded, when the wind is light, in a continuous cloud of smoke. This will, in many instances, prevent frost from occurring. The stakes must not be set so close to the fruit trees that the heat will affect the branches.

* * * * * * *

In France frost bells are sounded when frost is expected, and grape growers protect their crops by pouring tar on the ground among the vines and setting it on fire. The smoke thus made often saves the crop of an entire vineyard. Bituminous coal burned in baskets is less expensive and has been successfully used. Forty to fifty baskets of coal per acre are required, and the more used the better would be the result. Strawberry crops can be protected in this manner at a cost of $4 to $6 an acre.

* * * * * * *

The following letter from Capt. B. F. Johnson, president Gulf Coast Horticultural Association, gives an idea as to what can be done in protecting crops [by showing what was done in February, 1899]:

"I was in Galveston, Tex., on the morning of February 11, 1899, and upon hearing that a freeze was expected I called at the United States Weather Bureau office, and was informed that the hardest freeze of the season was threatened. I immediately gave orders over the long-distance telephone to my employees at Arcadia to cover up all hotbeds and also all plants already transplanted. All the plants in hotbeds were carried through the freeze by protection and saved from injury. I had about 7,500 cabbage plants set out, and on account of my employees not going to work as directed we did not reach these until ice had formed over some of the plants. I took the first train out of Galveston, which was in the afternoon, and upon my arrival at Arcadia we continued protection, notwithstanding the fact that ice had already formed over the plants. We placed a handful of hay over the north side of about 1,000 cabbage, and at least 750 of these came through the freeze uninjured, while all

the unprotected were killed. The varieties protected were the Jersey-Wakefield and Henderson-Succession, and I could see no difference between the two after the freeze. Had my employees carried out the instructions given them in the morning I would have saved 75 per cent of the 6,500 unprotected plants which were killed, and this would have yielded me about $150 net. I had 1¾ acres of Irish potatoes planted in January, which were up to fair stand. These were not protected and were all killed. I had about one-sixteenth acre planted with hand planters the 1st of February. These were protected with vines from old sweet potatoes grown on the ground the previous year. Fully one-eighth of these came up after the freeze with this light protection, and thus enabled me to put the first potatoes on the Texas market, and they brought me $3 a bushel net. If I had been prepared and protected the 1¾ acres planted in January I would have received a nice little sum, and more than ten times the cost of protection. Strawberries not mulched were virtually killed, while those mulched made an average crop. The great value of protection to berries in case of freezing weather has been forcibly illustrated. The entire crop of early berries (the only berry crop which pays well) would have been a total failure without protection."

* * * * * * * *

The following communication of Mr. M. M. Cox, president of the League City Truck Growers' and Farmers' Association, League City, Tex., gives further the manner and value of protecting crops from the frost and freeze during the abnormally cold weather of February 12 and 13, 1899. Mr. Cox says:

"I had two acres of strawberries—Noonan and Mitchell varieties—which were blooming and fruiting nicely. In anticipation of a freeze we placed hay for mulching purposes between the rows, which were about 3 feet apart. Upon the receipt of the telegram, Saturday morning, February 11, 1899, announcing the approach of a freeze, I got word to the man who had charge of my berry crop. He immediately went to covering the vines with hay previously placed in the berry patch, and succeeded in covering 20 rows. The vines of the protected berries were practically uninjured, while the unprotected plants were frozen to the ground and many of them totally killed. From these 20 rows, about one-fourth acre of protected plants, we picked berries nearly three weeks before the unprotected berries were ready for market. The one-fourth acre of protected berries yielded me more than the remaining 1¾ acres altogether. In fact, we got the bulk of our crop—a little more than $300 net—off this one-fourth acre. If we had used more hay for covering we could have saved the blooms and berries already on the vines notwithstanding the severity of the freeze, as some berries were saved as it was.

"Others in League City protected berry crops with the same results as those which attended our protection.

"Hotbeds protected for ordinary freezes were killed, but some put extra covering on the beds, using blankets, and thus saved their plants without injury. Those who protected in this manner were setting their plants in the field while others were sowing seeds in the hotbeds, and were thus at least four weeks ahead with their crops for the market.

"The warnings of the Weather Service promise much for the coast country."

Mrs. C. W. Benson, Alvin, Tex., in writing of protection of crops, says:

"In past seasons the advance notices of frost and freezing weather given out by the United States Weather Bureau have been distributed among the truck growers by messenger service, and by this means have saved our planters who protect their crops thousands of dollars.

In 1899 all crops were carried through safe until the severe freeze of February 11, and in several instances tender crops were successfully protected and carried through this freeze even when the mercury stood at 4 degrees above zero or 28 degrees below the freezing point. The principal crops protected were tomatoes, cucumbers, cabbage, and strawberries. The three first mentioned were still in hotbeds or cold frames, and were protected by covering with hay to a depth of about 1 foot, over glass or

canvas. Berries were protected by turning over mulch and covering up plants from 4 to 6 inches in depth. This mulch had already been placed between the rows to retain moisture and keep the berries clean on plants. This protection did not save blooms, but the plants so protected came through the freeze with much less damage and produced more fruit than the unprotected plants. I judge that the protection was fully worth $30 per acre, as prices of berries remained firm all through the season on account of general scarcity of fruit. About 25 acres were thus protected from the freeze.

"The value of the tomato crop saved through protection from this freeze averaged fully $125 per acre in spite of the subsequent drought, which extended from February 14 until after the crop was marketed, with the exception of a few slight showers, which were insufficient for general growth of plants. With rain as in an average season the crop would have paid $250 per acre, and enough plants were carried through this freeze by protection to set 17 acres.

"About 200,000 cabbage plants were carried through this severe freeze of February and successfully transplanted. These would have netted the growers $8,000 with sufficient rain during the growing period. The value of the cucumber crop from actual results was about $1,000, which would have increased 1,000 per cent in an ordinary season, but they were held back for lack of moisture until the north Texas crop was on the market, as cucumbers which sold at from 40 to 50 cents per one-third bushel box would have netted from 75 cents to $1.25 per dozen.

"Protection of crops enables the grower to generally reach the first markets, but without the advance notices of frost and freezing weather it would be impracticable to protect; our people would bear the expense of the warnings rather than do without them."

WORK IN LOUISIANA AND FLORIDA.

Probably the most extensive experiments for the protection of citrus fruits in the Southern States have been conducted in a large orange orchard near Diamond, La., under the management of Mr. Hewitt Chapman. In the Report of the Louisiana Section of the Climate and Crop Service of the Weather Bureau for February, 1899, this grove and some of the methods employed to protect its crop are described by Mr. Alexander G. McAdie, of the Weather Bureau, as follows:

The orange orchard is about 43 miles south of New Orleans, near the river, and about 20 miles north of what is considered the safe orange belt of Louisiana. The grove covers 130 acres and contains 15,000 trees, including Creole Sweets, Mandarins, and Satsumas; sweets budded on Trifoliata and Pomeloes on sour stock. These young trees withstood the frosts of January 1, December 4, 9, 10, 11, 1898. The writer visited this grove on December 11, when ice one-quarter inch in thickness could be found along the roads and in exposed places. The orchard is irrigated by means of river water carried over the levee by a Knowles direct-acting pump, 6 by 8, worked by a 40-horsepower boiler. The pump dispenses with belt, pulley, and engine. To flood the whole orchard of 50 acres to a depth of 1 inch would require 5,650 tons of water, or 1,357,700 gallons. Barely one-hundredth part of this quantity of water is required. The water pumped from the river over the levee goes into a main ditch, from which run subditches, each subditch feeding two small inlets through the orchard between each line of trees. The distance between the trees averages 12 feet. The middle ground is cultivated and level so that the water can run in a steady stream to and out through the quarter drain into the drainage ditches which are 1 acre apart, running parallel into the rear canal.

The grove has a northern exposure; fronts on the river; the land runs west and east, falling slightly from the river; and there are no wind breaks. It is also proposed at this orchard to experiment with the various types of smudges. One portion, about 10 acres in extent, will be protected by smudge fires, some portable and

some stationary. In another section, covering about 30 acres, the bodies and lower portions of the orange trees are to be wrapped with native hay. In the third section the dirt is to be hilled up about the trees; while in the fourth the trees are to be wrapped with floor matting.

In the methods described above for the protection of trees it was thought that colder weather than that of February, 1895, was not likely to occur again. It appears that the pumping plant, on which great reliance was placed, did successfully hold back the trees up to about February 2, 1899, but the extremely warm weather of the 3d, 4th, and 5th started the sap and forced growth. It is doubtful, however, whether water protection is adequate under such severe tests. All the orange trees above ground appear to have suffered badly, in nearly every case the tree being killed. Those trees which were wrapped with matting likewise suffered and the protection again appears to have been inadequate to withstand such extreme cold. In the section where the earth was hilled up around the trees the results were very satisfactory. It is thought that about 4,000 trees have been saved by this means. An inspection of the orchard as late as March 5 indicates that notwithstanding the severity of the test, there is reason to believe that orange trees can be protected successfully by hilling up the earth.

The devices employed in Louisiana, as described above, were found to be an effective protection against damage by frost, and the protection afforded by irrigation and by wrapping the bodies and lower portions of trees with hay was sufficient for ordinary freezes. During the unprecedented cold of February, 1899, the section in which dirt was hilled up about the trees was the only one in which trees escaped serious injury.

The devices for protection from frost used in Florida conform in character to those tested in the California experiments, in which coal fires were found to be the most effective means of protection, and appliances for adding moisture to the air were successful only to a degree. As artificial appliances are totally inadequate to add to the atmosphere any very appreciable amount of moisture, it is evident that methods which may have been found ineffective in the dry climate of California would possess value in locations where their office is confined to adding moisture to an already moist atmosphere. The Gulf and South Atlantic Coast States, and in fact the country generally from the Mississippi Valley to the Atlantic seaboard possesses a moist atmosphere, and the fruits and tender garden vegetables of these districts can, therefore, be the more readily protected from frost by devices which add moisture to the air. As a means of adding moisture to the air, irrigation should be more effective than in California, and in localities where this method can be used, protection would be assured except against hard freezes.

Owing to the comparatively inexpensive character of the materials used in damp smudge fires, they seem the best adapted for common use in orchards, vineyards, and gardens. Berries and other low plants can be protected with but little expense by coverings of straw and other light materials. Devices for actually heating the free and open air are expensive and of doubtful utility, and their value is dependent solely upon a comparatively still air and small, numerous, and well-distributed fires.

In Florida many experiments have been made with a view to adopting devices which will protect orange trees from injury during the periods of severe cold which at times visit that section. The fact that these periods of cold are infrequent does not relieve the grower from the necessity of providing for their occurrence. The most economic of the devices tested appears to be the "banking up" of trees with dirt or sand. In Florida, as in Louisiana, this has been found to be a very effective means of protection. The trunks of the older trees can be banked, and the younger trees can be almost entirely covered without necessarily causing any damage to the trees.

The more expensive methods include warming the air by means of open wood fires and sheet-iron stoves. These methods have been found to be fairly satisfactory on still nights. Coverings of cloth and other suitable materials are stretched on frames and the confined air is warmed by means of stoves. One extensive orange grower is building board sheds over his groves, the interiors of which will be heated by wood fires. When danger from frosts and freezes is past, the tops and sides of the sheds can be removed. On account of their great cost, coverings of this character can not, of course, be generally used.

FROST SEASONS FOR SEVERAL CROPS.

Damaging frost is likely to occur in Florida from the middle of October until nearly the middle of April. During this period pineapples which are not in the lee of bodies of water should be afforded means of protection. Strawberries are subject to damage from January to the early part of April; oranges from December to the middle of February, and orange trees are subject to damage by freezes during December, January, and February. Garden truck is especially subject to injury from January 1 to April 15. In the trucking districts of the interior of the Gulf and South Atlantic States the period during which damaging frost is likely to occur extends from October to April, inclusive.

Temperature conditions are of great importance in the culture of rice, and a severe frost late in October may injure the grain so much as to cause its discoloration. Owing to the manner of growth and culture of this crop and the character and extent of the fields it covers, practically nothing can be done to avert injury from heavy frosts that may occur about the time of harvest.

Tobacco growers may protect their plant beds by coverings, but during the critical periods following transplanting and approaching harvest severe frost will ruin plants in fields which are too large for the successful application of the devices herein described. During the maturing season tobacco plants in large fields can be given protection from damage by frost only by cutting and "frost heaping" on the eve of the occurrence of frost.

Cotton will be seriously injured by low temperature early in the spring whether frost occurs or not. Owing to the area covered by the

fields, no adequate means of protection against cold and frost, suitable for general use, has been discovered.

In Louisiana sugar cane is planted in the fall, winter, and early spring. The plants are often killed or seriously injured by frosts in the spring and in October and November. When frost is expected, the cane can be cut close to the ground and "mattressed," the product of three rows being so laid together that the leaves of one armful will cover the butts of the preceding; being thus thatched the canes are protected from frost and will keep in this state for several weeks without injury.

In the cranberry districts of New Jersey, Cape Cod, and Wisconsin, and in limited areas in other parts of southeastern New England, and the upper lake districts, which are devoted to cranberry raising, flooding is the only effective means of protection against damage by frost. This crop is confined to swamp or bog lands, and when danger from frost is apprehended, appliances for letting in a sufficient quantity of water to keep comparatively warm and moist the lower stratum of air are usually provided.

Young potato plants can be saved from damage by frost by a covering of dirt. In large patches this can be accomplished by plowing a furrow alongside the rows of plants and allowing the dirt to bury them.

Gardens can be protected from frost by coverings of straw, paper, or other loose and light materials, or by damp smudges; and the latter device appears to be, in point of effectiveness and cost, the best adapted for general use not only for small areas devoted to gardens, but also for orchards, vineyards, and fields. The effectiveness of any method of protection is of course decreased as the area for which protection is desired is increased, and there is no process yet devised that can give adequate protection against frost and cold in the great grain fields of the country.

CONCLUSIONS.

Experiments and observation seem to establish the following facts: The danger of damage from frost can be materially lessened by placing early and tender plants on high grounds and crests, and hardier plants in low grounds and hollows. When ground can be selected in the lee, or to the south and east of considerable bodies of water, the danger will be further lessened.

In the dry climate of the citrus fruit region of California and in the promising fruit districts of Arizona small and numerous fires, preferably of coal burned in iron baskets, have been found to be the most effective device used for protection against frost; second in point of utility may be placed irrigation; and the practicable process which affords the least protection in that district appears to have been smudge fires. As before stated, the unsatisfactory character of the experiments made with smudge fires may be attributed to the fact that artificial devices are inadequate to charge dry air with sufficient moisture to prevent loss of heat from the earth by radiation. Reports of California

experiments do not distinguish between frosts and freezes. Owing, however, to the usually moderate character of the cold periods of that section the devices referred to would doubtless be equally effective against frosts and freezes.

In the orange-growing districts of the South irrigation affords the most effective protection against frost, while in sections where this process can not be employed damp smudge fires properly handled are best adapted to general use. Severe freezes which kill or injure the trees can be guarded against only by housing the trees and heating the confined air, or by "banking up" the trees with dirt or sand. The latter method is comparatively inexpensive and is therefore the better adapted to general use.

In all sections loose coverings may be used to advantage for protecting low plants, and damp smudge fires may be profitably employed in gardens, orchards, and small fields.

It is important to remark in this connection that none of the artificial devices described should be employed during the growing season except on the immediate eve of threatened frost. The necessity for observing this precaution becomes apparent when the fact is considered that any appliance for protecting plants from cold will maintain abnormally high temperatures when average weather conditions prevail. The effect of this process would be to force the growth of the plants and render them the more susceptible to damage by the late frosts of spring. In all sections and localities reached by the forecasts and special warnings of the Weather Bureau action to put into operation protective measures can be safely deferred until warning notices are issued. The Bureau has made provision for a distribution of frost warnings which is limited only by the resources of the telegraph, telephone, mail, train, and signal services, and agriculturists whose location renders possible the receipt of the warnings by any of the means referred to should adopt measures for securing them. That action in this direction would be profitable is shown by the testimony of those who have received substantial benefits from the warnings.

FARMERS' BULLETINS.

These bulletins are sent free of charge to any address upon application to the Secretary of Agriculture, Washington, D. C. Only the following are available for distribution:

No. 16. Leguminous Plants for Green Manuring and for Feeding. No. 19. Important Insecticides: Directions for their Preparation and Use. No. 21. Barnyard Manure. No. 22. Feeding Farm Animals. No. 23. Foods: Nutritive Value and Cost. No. 24. Hog Cholera and Swine Plague. No. 25. Peanuts: Culture and Uses. No. 26. Sweet Potatoes: Culture and Uses. No. 27. Flax for Seed and Fiber. No. 28. Weeds; and How to Kill Them. No. 29. Souring of Milk and Other Changes in Milk Products. No. 30. Grape Diseases on the Pacific Coast. No. 31. Alfalfa, or Lucern. No. 32. Silos and Silage. No. 33. Peach Growing for Market. No. 34. Meats: Composition and Cooking. No. 35. Potato Culture. No. 36. Cotton Seed and Its Products. No. 37. Kafir Corn: Characteristics, Culture, and Uses. No. 38. Spraying for Fruit Diseases. No. 39. Onion Culture. No. 40. Farm Drainage. No. 41. Fowls: Care and Feeding. No. 42. Facts About Milk. No. 43. Sewage Disposal on the Farm. No. 44. Commercial Fertilizers. No. 45. Some Insects Injurious to Stored Grain. No. 46. Irrigation in Humid Climates. No. 47. Insects Affecting the Cotton Plant. No. 48. The Manuring of Cotton. No. 49. Sheep Feeding. No. 50. Sorghum as a Forage Crop. No. 51. Standard Varieties of Chickens. No. 52. The Sugar Beet. No. 53. How to Grow Mushrooms. No. 54. Some Common Birds in Their Relation to Agriculture. No. 55. The Dairy Herd: Its Formation and Management. No. 56. Experiment Station Work—I. No. 57. Butter Making on the Farm. No. 58. The Soy Bean as a Forage Crop. No. 59. Bee Keeping. No. 60. Methods of Curing Tobacco. No. 61. Asparagus Culture. No. 62. Marketing Farm Produce. No. 63. Care of Milk on the Farm. No. 64. Ducks and Geese. No. 65. Experiment Station Work—II. No. 66. Meadows and Pastures. No. 67. Forestry for Farmers. No. 68. The Black Rot of the Cabbage. No. 69. Experiment Station Work—III. No. 70. The Principal Insect Enemies of the Grape. No. 71. Some Essentials of Beef Production. No. 72. Cattle Ranges of the Southwest. No. 73. Experiment Station Work—IV. No. 74. Milk as Food. No. 75. The Grain Smuts. No. 76. Tomato Growing. No. 77. The Liming of Soils. No. 78. Experiment Station Work—V. No. 79. Experiment Station Work—VI. No. 80. The Peach Twig-borer—an Important Enemy of Stone Fruits. No. 81. Corn Culture in the South. No. 82. The Culture of Tobacco. No. 83. Tobacco Soils. No. 84. Experiment Station Work—VII. No. 85. Fish as Food. No. 86. Thirty Poisonous Plants. No. 87. Experiment Station Work—VIII. No. 88. Alkali Lands. No. 89. Cowpeas. No. 90. The Manufacture of Sorghum Sirup. No. 91. Potato Diseases and Their Treatment. No. 92. Experiment Station Work—IX. No. 93. Sugar as Food. No. 94. The Vegetable Garden. No. 95. Good Roads for Farmers. No. 96. Raising Sheep for Mutton. No. 97. Experiment Station Work—X. No. 98. Suggestions to Southern Farmers. No. 99. Common Insects on Shade Trees. No. 100. Hog Raising in the South. No. 101. Millets. No. 102. Southern Forage Plants. No. 103. Experiment Station Work—XI. (In press.)

O

Cop. 3

Issued February 14, 1908.

1
Ag84F
Reserve

U. S. DEPARTMENT OF AGRICULTURE.

FARMERS' BULLETIN 104

Rev.ed.
follows

NOTES ON FROST.

BY

E. B. GARRIOTT,
Professor of Meteorology, Weather Bureau.

Revised January, 1908

WASHINGTON:
GOVERNMENT PRINTING OFFICE
1908.

348256

LETTER OF TRANSMITTAL.

U. S. DEPARTMENT OF AGRICULTURE,
OFFICE OF THE WEATHER BUREAU,
Washington, D. C., January 25, 1908.

SIR: I have the honor to recommend that the manuscript herewith be published as a revised edition of Farmers' Bulletin No. 104.

In compliance with directions contained in your letter of March 29, 1899, a paper on the subject of frost, prepared by Prof. E. B. Garriott, Weather Bureau, was submitted and published as Farmers' Bulletin No. 104. This bulletin furnished valuable information regarding methods of protection from frost. Methods advocated in previous publications were mainly the results of experiments conducted in California orchards. Bulletin No. 104 was designed to be of value in all sections of the country rather than in restricted districts.

As the California experiments quoted were perhaps the most thorough ever made in this country, a description of the methods employed and the results obtained is reprinted in brief in this paper, together with results of similar experiments made in the citrus-fruit and truck-growing districts of the Gulf States, where varying climatic conditions have been found to produce somewhat different results. Additional information personally acquired by the author during a visit to Florida in the winter of 1906–7, and certain climatic data, also form features of this paper in its revised form.

Your wishes that the subject be treated in a popular way and divested of all technicalities, and made applicable to all portions of the United States, have been complied with so far as the nature of the subject will permit.

Respectfully,

WILLIS L. MOORE,
Chief, Weather Bureau.

HON. JAMES WILSON,
Secretary of Agriculture.

CONTENTS.

	Page.
How frost is formed	7
Seasons of frost	8
When to expect frost	9
Dates of early autumn and late spring frosts	13
Protection from frost	15
Devices for preventing rapid radiation of heat	16
Glass screens	16
Cloth screens	16
Lath screens	16
Other methods	16
Smudge fires	16
Devices for adding moisture to the air	17
Fires of damp straw and stable manure	18
Sacks of manure	18
Bales of wet straw	18
Prunings	18
Portable smudge fires	18
Devices for charging the air with moisture	19
General observations and practical results	22
Work in Texas	22
Work in Louisiana	25
Work in Florida	26
Frost seasons for several crops	28
Conclusions	30

104

NOTES ON FROST.

HOW FROST IS FORMED.

The atmosphere of the earth always contains more or less moisture in an invisible form. When at a considerable elevation above the earth this moisture, or aqueous vapor, is condensed, clouds are formed; when the process of condensation is more active and the temperature of the air is above freezing, rain falls; and when the temperature of the air is below freezing, snow is produced. When the moisture of the air in immediate contact with the earth is condensed at temperatures above freezing, dew is formed; when at temperatures below freezing, frost is deposited. Frost is, therefore, the moisture of the air condensed at freezing temperatures upon plants and other objects near the surface of the earth.

In the process of frost formation the temperature of the air a few feet above the earth is commonly several degrees above freezing. The surfaces upon which frost is deposited must, however, possess freezing temperatures. The manner in which frost is deposited on plants and other objects is very similar to that in which the air moisture of a room is frozen and deposited upon window glass, the temperature of which has been reduced to freezing by the out-of-doors cold. In the case of the frost on the window glass the process is one that can readily be understood. Some explanation, however, is required of the formation of frost, which requires freezing temperature, when at times a temperature above freezing is registered a few feet above the surfaces upon which frost appears.

There are several processes by which the temperature of plants may be reduced below the temperature of the air which surrounds them. The most important of these processes is radiation, by means of which heat escapes from objects and passes into the surrounding air.

In the frost-forming process heat from the sun which is absorbed by the earth and by plants during the day is lost by radiation at night. During the day the earth both absorbs and reflects the heat received from the rays of the sun, and the lower stratum of air is warmed by this reflected heat. During the night when no direct heat is received from the sun's rays the lower air stratum receives no reflected heat,

and, at the same time, heat which has been absorbed by the earth is radiated, or, in other words, it rises through the overlying air, though not necessarily and entirely as sensible heat; that is, heat perceptible to the senses. In the frost-forming process there is another very important factor in the production of cold; that is, the evaporation of moisture from the earth and from plants. The nature and composition of frost require that the air in immediate contact with the surfaces upon which frost forms shall contain moisture. This is moisture that has been stored in the earth in visible form, i. e., in the form of water, and which has escaped from the earth, not as visible moisture, but by the process of evaporation and in the form of aqueous vapor. Some part of the heat stored by the earth during the day is, therefore, lost in evaporating the moisture contained in plants and in the earth. The heat utilized in this process is retained in the air, not as sensible, but as latent heat. Heat is rendered latent, in the same manner, when it is used to boil away water, and it again becomes sensible or liberated heat when the water vapor produced by boiling, which is a rapid process of evaporation, is again condensed into water. It appears, therefore, that while the temperature of surfaces upon which frost forms, and of the air in immediate contact with them, is lowered by the evaporation of moisture from the surfaces, the influence of the process does not extend to air a few feet above the ground.

Another method by which plants lose their heat is by convection, by means of which they are chilled by contact with colder air. This process, while important in the presence of freezing-air temperatures, can scarcely be considered a factor in the formation of frost proper, which is usually accomplished when temperature observations show the air to be above freezing.

Another element in the formation of frost is found in the fact that air, like fluids, arranges itself according to its density or weight. Thus the air in immediate contact with the earth becomes heavier as its temperature is lowered by the radiation of heat and settles into depressions or valleys and over lowlands, causing, at times, frost in low-lying districts, while neighboring higher grounds escape the visitations.

SEASONS OF FROST.

Generally speaking, agricultural products in some part of the United States are menaced by frost from the time nature begins to extend her mantle of green from the citrus regions of the South toward the grain fields of the North until the verdured area contracts southward before the first winds of winter; from the hour that seeds germinate and buds blossom until crops are gathered; from early spring until late fall. As a rule, the critical periods are during the early growth of plants and when the fruit is in the bloom, and again

when crops are maturing. In the first instance vegetation is threatened by the frosts of early spring, and in the latter by the frosts of autumn.

Considering the several agricultural sections in greater detail we find that damaging frost is likely to occur in the north half of the Florida peninsula and in the region immediately bordering the coast of the Gulf of Mexico, and in the Pacific Coast States from the latter part of October until the early part of April, and in the North Pacific Coast States as early as the middle of October and as late as the last week of April. From December until early in March the frost limit and even the line of freezing temperature may, at long intervals, be extended southward over central and southern Florida. The period of damaging frosts in the interior of the South Atlantic and Gulf States extends from November to April. From October to April the region in which agricultural products are subject to damage by frost is extended to the southern lake region, over the Upper Mississippi and Lower Missouri valleys, Kansas, and Nebraska, and from September to May frost visitations are usually confined to the extreme Upper Mississippi, Upper Missouri, and Red River of the North valleys, and the Rocky Mountain and plateau regions from central New Mexico and the Texas Panhandle northward.

The occurrence of summer frosts in mountain regions and in extreme northern districts of the country is common. The following are notable instances of summer frosts in agricultural districts:

August 22 to 24, 1890, a frost-bearing cool wave advanced from the Dakotas over the lake region and the interior, or mountain districts, of the middle Atlantic States.

August 28 to 31, 1892, a cool wave advanced from the Rocky Mountain and plateau regions, with frost from New Mexico to the Dakotas, Iowa, Minnesota, northern Wisconsin, and northern Michigan.

August 8, 1904, frost damaged vegetation in low-lying lands in the States of the upper lake region.

Bulletin P, of the Weather Bureau, gives a chronological statement of the more important cold periods that have been experienced in the United States, and contains charts that present conditions that have preceded and attended the principal cold waves and frosts of the last fifteen years.

WHEN TO EXPECT FROST.

As the conditions which produce damaging frost are subject to modifications which are as numerous as the kinds of crops raised, and as varied as local topography, local climate, and local soil conditions in the various sections of the United States, the writer in attempting to treat briefly a subject which is so diversified in its aspects is confronted with many difficulties. Suffice it to say that with other atmos-

pheric conditions favorable for its occurrence frost may, as a rule, be expected when temperature, as reported by stations of the Weather Bureau, falls to a point 8° to 10° above the freezing point. As previously stated, the surfaces upon which frost is deposited must possess freezing temperature, although the temperature of the air a few feet above the surfaces may be several degrees above freezing; and it is the temperature of the air, in some instances many feet above the ground, that is given by the Weather Bureau observations. Another atmospheric condition favorable for the occurrence of frost is a clear, cloudless, and comparatively calm night. The presence of clouds retards radiation or loss of heat from plants; the clouds act as a screen in preventing the heat collected from the sun's rays during the day from escaping into the upper air. When clouds are not present and a withdrawal of the sun's rays causes a rapid cooling of the air at moderate elevations, the warmer air which collects near the surface of the earth during the day rises, and the cooler upper air, owing to its greater density or weight, settles to the earth. It will be noted that clouds not only prevent the escape of the warm air in immediate contact with the earth, but also blanket, as it were, the upper part of the lower air stratum. Calm or comparatively still air is a condition which favors the formation of frost. On windy nights the air is disturbed and is not permitted to arrange itself in layers according to its density, with the densest and coldest air near the surface of the earth; it is kept mixed up by the wind.

A very interesting fact, which illustrates the manner in which cold air settles by its greater weight to the surface of the earth and allows the warmer air to assume higher levels, causing frost in low grounds and granting to the higher grounds a comparative immunity from frost, is set forth in the following article, by Prof. Cleveland Abbe, in the Monthly Weather Review for December, 1893, entitled Thermal Belts, Frostless Belts, or Verdant Zones:

These are local names given to certain regions on mountain sides within which nocturnal frost rarely or never occurs in the springtime, although freezing temperatures occur in the winter time, consequently tender vegetation flourishes with remarkable vigor in these regions. The following are the only references to thermal belts in the United States at present known to the editor, but as such regions are specially important to the horticulturist and agriculturist it is hoped that the correspondents of the Weather Bureau will bring these thermal belts to notice wherever they occur in order that their meteorological peculiarities may be better understood.

In the Agricultural Report of the Patent Office for 1861 Mr. Silas McDowell, of Franklin, Macon County, N. C., describes the verdant zone in that county in the valley of the Little Tennessee River. It occupies the region between 300 and 700 feet above the valley of the river, which latter is about 2,000 feet above sea level. On tracing this zone up among the smaller tributaries of the Tennessee River he found that in the higher valleys, where the bottom land is about 3,900 feet above sea level, the verdant zone lies between 4,000 and

4,100 feet. Within this zone frost never injures the vegetation, and the most tender grapes never fail to produce abundant crops.

Prof. J. W. Chickering, jr., in the bulletin of the Philosophical Society of Washington, March, 1883, and in the American Meteorological Journal, vol. 1, describes the following thermal belt: "In Polk County, N. C., along the eastern slope of the Tryon Mountain range, in latitude north 35, the thermal belt begins at the base of the mountain, at an elevation of 1,200 feet above the sea, and extends up 2,200 feet, being most perfect at about 1,500 feet. It is about 8 miles long, and is distinguished by magnificent flora, such as would be characteristic of a point 3 degrees south of the actual latitude."

Prof. John Leconte, of Berkeley, Cal., in Science, vol. 1, p. 278, states that at Flat Rock, near Hendersonville, Henderson County, N. C., on the flank of the mountain spur adjacent to the valleys of the Blue Ridge, he also observed a frostless zone. The valley is about 2,200 feet above sea level, and the thermal belt is 200 to 300 feet above the valley.

Mr. J. W. Pike, of Vineland, N. J., states that among the mountains of California he has discovered that during the night the cold is much greater in the valleys than on the terraces several hundred feet above, due to the settling of the cold air, so that a thermal belt is formed at that height separating the frosty valleys from the colder highlands.

In the Tennessee Journal of Meteorology for January, 1894, published by the State weather service, the author describes a thermal belt between Los Angeles and the Pacific coast. It traverses the foothills of the Cahuenga range, and has an elevation of between 200 and 400 feet and a breadth of about 3 miles. It occupies the midway region of the range.

In the American Meteorological Journal, vol. 1, Mr. S. Alexander describes a thermal belt in which the peach tree flourishes in the southeastern portion of Michigan. He shows that the cold island discovered by Winchell in that region is really the bottom of a topographical depression into which the cold air settles. It is a long valley surrounded by a belt of elevated country from 50 to 600 feet above lakes Michigan and Huron. The valley and the isotherms trend northeast and southwest from Huron County through Sanilac, Lapeer, Oakland, Livingston, and Washtenaw to Hillsdale County. The highlands of this region are all much freer from frost than the lowlands, and all much more favorable for early vegetation. He does not state that any point is high enough to be above the thermal belt, but that, in general, two equal parallel thermal belts inclose the cold island between them.

It is generally conceded that these thermal belts depend both upon the drainage of cold air downward into the lower valleys and the freedom of radiation from the surface of the ground to the clear sky overhead. During a still night, when frosts occur, the surface of the hillside cools by radiation, and hence cools the air in contact with it; the latter flows downward as long as its cooling by radiation and conduction exceeds its warming by compression. Inasmuch as its cooling depends on contact with a still colder soil or plant, it soon accumulates in the lowlands as a layer of cold air, which grows thicker during the night by the steady addition of the thin layer of descending air in contact with the ground on the hillsides. The warmer air, which has not yet had an opportunity to cool by contact with the ground, floats on top of the cold mass; it spreads out toward the hills, and is continuously furnishing its heat to the adjacent hillsides as fast as it comes in contact with them before it also cools and descends. The formation of the thermal belt seems to depend largely upon this gentle circulation during the nighttime. The lower limit of the belt is defined by the depth of the accumulation of cold air in the confined valley and rises higher in proportion as the night is clearer and longer, and also in propor-

tion as the valley is more or less perfectly inclosed. The upper limit of the thermal belt may depend upon the strength of the wind and the general temperature of the air. But if there be no wind, then it depends equally on the freedom of radiation to the clear sky and on the above-described circulation of air.

The facts quoted in this article show that frost may be expected on low grounds at times when higher grounds escape the visitations, and the lesson they teach is that early and tender crops should, so far as may be practicable, be confined to crests, hillsides, and mountain sides, and later and hardier crops to the lowlands and valleys.

Local climate, as it is influenced and regulated by the proximity of bodies of water, must be given great weight in calculations regarding frost. Frost is less likely to occur in localities swept by moisture-laden air which has crossed a considerable body of water. This is more especially the case in the fall of the year, when the temperature of bodies of water is reduced very slowly; and in the South, where the water temperature continues relatively high throughout the year, the influence of the water is especially marked. During the colder months air is not only warmed in crossing considerable bodies of water, but it also absorbs moisture, which, although invisible in the form of water vapor, has the effect of retarding the radiation of heat from the earth. In fact, so pronounced is the influence of water and water vapor that localities which with reference to exposure to west and northwest winds are, as it were, in the lee of large bodies of water are comparatively free from frost visitations. On the keys of the east Florida coast which are protected on the west by lagoons, rivers, lakes or bays, frost is said to be unknown, and throughout the central and northern sections of the United States, localities which are protected on the west and northwest by bodies of water of considerable size are not only favored, by reason of their position, with conditions which do not promote the formation of frost, but also receive the benefits derived from heavier falls of snow.

Local soil conditions constitute, to a certain extent, a factor in the formation of frost. As a rapid loss of heat is promoted by an active evaporation of moisture, it is evident that, with other conditions equal, frost is more likely to occur on damp than on dry ground, provided, however, that the ground is not too moist, for in the latter event the amount of moisture evaporated and added to the air would have a tendency to retard the radiation of heat from the earth. And herein lies the distinction, so far as frost formation is concerned, between moist air and moist soil. An excess of moisture in the air, in preventing, to a degree, the radiation of heat, is unfavorable to the formation of frost. As frost is the moisture of the air in immediate contact with the earth condensed at freezing temperature, it follows that the earth, from which the moisture of the air is drawn, must contain more or

less water, and it is evident that damaging frost will occur with a limited rather than with a large quantity of moisture arising from the earth and a moderately moist and comparatively still air. Both of these conditions usually obtain following, but not too closely following, the rains of spring and autumn. The character of the soil as regards its capacity for retaining moisture must also be taken into account, more especially during periods of comparatively dry weather. Moist soil, and the vegetation which springs from it, is chilled to a lower temperature by the evaporation of moisture, and is therefore more subject to visitations of frost than soil which is dry, or into which water penetrates deeply, leaving the surface dry, or soil which sheds the rain and does not absorb and hold water. It is important to note, however, that very moist soil, or soil which contains a large amount of surface water, is, owing to the quantity of moisture yielded to the air, not so subject to the heavy and damaging frosts which visit plants on moderately damp ground or ground that has a small capacity for moisture.

DATES OF EARLY AUTUMN AND LATE SPRING FROSTS.

The following table shows for stations of the Weather Bureau throughout the United States dates of first killing frost in autumn, and of last killing frost in spring; also the average dates of first killing frost in autumn and of last killing frost in spring:

Dates of first killing frost in autumn and of last killing frost in spring at Weather Bureau stations throughout the United States.

Station.	Date of earliest recorded killing frost in autumn.	Date of latest recorded killing frost in spring.	Average date of first killing frost in autumn.	Average date of last killing frost in spring.
Eastport, Me.	Sept. 3	June 19	Oct. 12	Apr. 28
Northfield, Vt.	Aug. 27	June 7	Sept.16	May 13
Portland, Me.	Sept.11	May 31	Oct. 18	Apr. 14
Boston, Mass.	Sept.30	May 11	Oct. 22	Apr. 26
Nantucket, Mass.	Oct. 1	Apr. 24	Nov. 5	Apr. 10
Block Island, R. I.	Oct. 30	May 11	Nov.16	Apr. 18
Narragansett, R. I.	Sept.21	May 30	Oct. 16	Do.
New Haven, Conn.	Sept.15	----do----	Oct. 17	Apr. 20
Binghamton, N. Y.	Sept.16	May 21	Oct. 6	Apr. 27
Albany, N. Y.	Sept.23	May 30	Oct. 17	Apr. 24
New York, N. Y.	Oct. 15	Apr. 30	Nov. 6	Apr. 10
Harrisburg, Pa.	Oct. 3	Apr. 26	Oct. 24	Do.
Philadelphia, Pa.	----do----	Apr. 29	Oct. 30	Apr. 8
Atlantic City, N. J.	Oct. 1	Apr. 25	Nov. 4	Apr. 11
Baltimore, Md.	Oct. 6	May 3	----do----	Apr. 4
Washington, D. C.	Oct. 2	May 11	Oct. 21	Apr. 7
Lynchburg, Va.	Oct. 4	May 7	Nov. 1	Apr. 14
Norfolk, Va.	Oct. 15	Apr. 26	Nov.12	Mar. 27
Wytheville, Va.	Sept.14	May 26	Oct. 7	Apr. 23
Charlotte, N. C.	Oct. 8	Apr. 26	Nov. 4	Apr. 1
Raleigh, N. C.	----do----	May 6	----do----	Apr. 5
Hatteras, N. C.	Nov. 7	Apr. 19	Dec. 11	Feb. 28
Wilmington, N. C.	Oct. 16	May 1	Nov.15	Mar. 27
Charleston, S. C.	Nov. 9	Apr. 2	Nov.30	Mar. 1
Columbia, S. C.	Oct. 19	Apr. 17	Nov. 8	Mar. 23
Augusta, Ga.	Oct. 8	----do----	Nov. 9	Mar. 20
Savannah, Ga.	Nov. 1	Apr. 5	Nov.26	Feb. 27
Jacksonville, Fla.	Nov.12	Apr. 6	Dec. 6	Feb. 19
Jupiter, Fla.	Nov.18	Apr. 7	Dec. 29	Feb. 14

Dates of first killing frost in autumn and of last killing frost in spring at Weather Bureau stations throughout the United States—Continued.

Station.	Date of earliest recorded killing frost in autumn.	Date of latest recorded killing frost in spring.	Average date of first killing frost in autumn.	Average date of last killing frost in spring.
Atlanta, Ga.	Oct. 4	Apr. 10	Nov. 7	Mar. 24
Macon, Ga.	Nov. 5	Apr. 18	Nov. 14	Mar. 15
Tampa, Fla.	Nov. 28	Mar. 19	Jan. 9	Feb. 8
Pensacola, Fla.	Nov. 12	Apr. 6	Dec. 5	Feb. 23
Mobile, Ala.	Oct. 31	Mar. 28	Nov. 30	Feb. 24
Montgomery, Ala.	Oct. 21	Apr. 5	Nov. 8	Mar. 11
Meridian, Miss.	Oct. 8	Apr. 10	Nov. 2	Mar. 20
Vicksburg, Miss.	Oct. 19	Apr. 6	Nov. 12	Mar. 17
New Orleans, La.	Nov. 19	Mar. 27	Dec. 15	Jan. 24
Shreveport, La.	Oct. 20	Apr. 2	Nov. 11	Mar. 4
Fort Smith, Ark.	Oct. 15	Apr. 6	Nov. 5	Do.
Little Rock, Ark.	Oct. 22	Apr. 14	Nov. 10	Mar. 20
Palestine, Tex.	Oct. 20	May 30	Nov. 13	Mar. 13
Galveston, Tex.	Dec. 4	Mar. 1	Dec. 25	Feb. 5
San Antonio, Tex.	Nov. 9	Mar. 20	Nov. 30	Feb. 25
Corpus Christi, Tex.	Nov. 30	Mar. 19	Dec. 25	Feb. 27
Memphis, Tenn.	Oct. 2	Apr. 16	Oct. 28	Mar. 24
Nashville, Tenn.	Oct. 8	May 14	Oct. 24	Apr. 2
Chattanooga, Tenn.	Sept. 30	---do---	Oct. 26	Do.
Knoxville, Tenn.	Oct. 1	Apr. 24	Oct. 27	Apr. 3
Louisville, Ky.	Sept. 24	May 14	Oct. 29	Apr. 6
Lexington, Ky.	Sept. 30	May 20	Oct. 23	Apr. 18
Evansville, Ind.	Sept. 10	Apr. 21	Oct. 30	Apr. 10
Indianapolis, Ind.	Sept. 21	May 21	Oct. 19	Apr. 16
Cincinnati, Ohio.	Sept. 30	Apr. 24	Oct. 25	Apr. 14
Columbus, Ohio.	Sept. 21	May 17	Oct. 16	Apr. 16
Elkins, W. Va.	Sept. 28	May 24	Oct. 8	May 12
Parkersburg, W. Va.	Sept. 24	May 22	Oct. 17	Apr. 11
Pittsburg, Pa.	Sept. 25	May 29	Oct. 19	Apr. 26
Oswego, N. Y.	---do---	---do---	---do---	Apr. 27
Rochester, N. Y.	Sept. 26	May 30	---do---	May 1
Buffalo, N. Y.	Sept. 23	May 25	Oct. 16	Apr. 25
Erie, Pa.	Oct. 12	May 17	Oct. 19	Apr. 20
Cleveland, Ohio.	Oct. 2	May 22	Oct. 31	Apr. 16
Sandusky, Ohio.	Oct. 8	May 2	Oct. 26	Apr. 14
Toledo, Ohio.	Sept. 9	May 29	Oct. 15	Apr. 24
Detroit, Mich.	Sept. 21	May 31	Oct. 11	Apr. 30
Port Huron, Mich.	Sept. 22	June 6	Oct. 9	May 8
Alpena, Mich.	Sept. 6	June 9	Sept. 26	May 14
Sault Ste. Marie, Mich.	Sept. 5	May 29	Sept. 24	May 16
Marquette, Mich.	Aug. 22	June 11	Oct. 2	May 15
Escanaba, Mich.	Sept. 9	June 16	Oct. 1	May 14
Green Bay, Wis.	Sept. 16	May 30	Oct. 4	May 5
Grand Haven, Mich.	Sept. 23	May 28	Oct. 10	Apr. 28
Milwaukee, Wis.	Sept. 25	May 29	---do---	Apr. 29
Chicago, Ill.	Sept. 18	---do---	Oct. 15	Apr. 18
Duluth, Minn.	Sept. 15	June 8	Oct. 14	May 3
Minneapolis, Minn.	Sept. 13	May 20	Oct. 8	Apr. 27
St. Paul, Minn.	Sept. 20	May 25	Oct. 5	May 6
Madison, Wis.	Sept. 30	May 13	Oct. 17	Apr. 21
La Crosse, Wis.	Sept. 2	June 23	Oct. 8	May 2
Dubuque, Iowa.	Sept. 27	May 21	Oct. 12	Apr. 19
Davenport, Iowa.	Sept. 18	May 22	Oct. 14	Apr. 22
Des Moines, Iowa.	Sept. 12	---do---	Oct. 8	Apr. 28
Keokuk, Iowa.	Sept. 18	May 2	Oct. 17	Apr. 2
Hannibal, Mo.	Sept. 30	May 14	Oct. 15	Apr. 14
Springfield, Ill.	Sept. 25	May 22	Oct. 16	Apr. 20
St. Louis, Mo.	Sept. 30	---do---	Oct. 27	Apr. 2
Cairo, Ill.	---do---	Apr. 19	Oct. 28	Mar. 29
Springfield, Mo.	---do---	May 19	Oct. 13	Apr. 16
Columbia, Mo.	Sept. 28	May 6	Oct. 15	Apr. 14
Kansas City, Mo.	Sept. 30	Apr. 23	Oct. 22	Apr. 7
Topeka, Kans.	Sept. 28	May 19	Oct. 13	Apr. 8
Wichita, Kans.	Sept. 23	Apr. 30	Oct. 19	Apr. 6
Concordia, Kans.	Sept. 27	May 19	Oct. 14	Apr. 24
Omaha, Nebr.	Sept. 18	---do---	Oct. 12	Apr. 15
Lincoln, Nebr.	Sept. 12	May 7	Oct. 9	Apr. 18
Valentine, Nebr.	---do---	June 21	Sept. 18	May 9
Sioux City, Iowa.	Sept. 13	May 21	Sept. 23	May 1
Huron, S. Dak.	Aug. 23	June 8	Sept. 20	May 13
Pierre, S. Dak.	Sept. 12	May 19	Sept. 30	Apr. 30
Moorhead, Minn.	Aug. 25	June 8	Sept. 22	May 14
Bismarck, N. Dak.	Aug. 17	June 3	Sept. 15	May 15
Williston, N. Dak.	Aug. 18	June 10	Sept. 14	May 18
Havre, Mont.	Aug. 14	June 6	Sept. 18	May 11
Helena, Mont.	Sept. 5	June 9	Sept. 25	May 10

Dates of first killing frost in autumn and of last killing frost in spring at Weather Bureau stations throughout the United States—Continued.

Station.	Date of earliest recorded killing frost in autumn.	Date of latest recorded killing frost in spring.	Average date of first killing frost in autumn.	Average date of last killing frost in spring.
Rapid City, S. Dak.	Sept. 13	May 20	Sept. 20	May 1
Lander, Wyo.	Aug. 23	June 18	Sept. 11	May 19
Salt Lake City, Utah.	Sept. 22	---do---	Oct. 18	Apr. 23
Grand Junction, Colo.	Sept. 14	Apr. 30	Oct. 29	Apr. 11
Cheyenne, Wyo.	Aug. 29	June 11	Sept. 16	May 22
North Platte, Nebr.	Sept. 10	May 23	Sept. 28	May 1
Denver, Colo.	Sept. 12	June 6	Oct. 7	May 6
Amarillo, Tex.	Oct. 16	May 23	Nov. 1	Apr. 16
Pueblo, Colo.	Sept. 12	---do---	Oct. 5	Apr. 26
Dodge, Kans.	Sept. 23	May 11	Oct. 15	Apr. 17
Oklahoma, Okla.	Oct. 7	Apr. 30	Oct. 31	Apr. 2
Fort Worth, Tex.	Oct. 22	May 1	Nov. 19	Mar. 18
Abilene, Tex.	Oct. 25	Apr. 16	Nov. 15	Mar. 15
El Paso, Tex.	Oct. 30	Apr. 22	Nov. 10	Mar. 20
Santa Fe, N. Mex.	Sept. 28	May 18	Oct. 19	Apr. 15
Yuma, Ariz.	None.	None.	None.	None.
Phoenix, Ariz.	Nov. 9	Mar. 31	Dec. 7	Feb. 18
Spokane, Wash.	Sept. 7	June 8	Oct. 17	Mar. 23
Walla Walla, Wash.	Sept. 28	May 3	Nov. 1	Apr. 6
Seattle, Wash.	Oct. 18	May 10	Nov. 22	Mar. 21
Portland, Oreg.	Oct. 13	May 9	Nov. 16	Mar. 17
Tatoosh Island, Wash.	Nov. 1	Apr. 19	Dec. 15	Mar. 14
Roseburg, Oreg.	Oct. 9	May 16	Oct. 30	Apr. 15
Baker City, Oreg.	Sept. 4	June 24	Sept. 27	June 3
Winnemucca, Nev.	Aug. 22	May 11	Sept. 24	June 20
Eureka, Cal.	Nov. 7	May 1	Dec. 8	Apr. 2
Red Bluff, Cal.	---do---	Apr. 9	Nov. 25	Mar. 15
Sacramento, Cal.	Oct. 28	Apr. 26	Nov. 15	Feb. 16
San Francisco, Cal.	Dec. 20	Apr. 20	Dec. 13	Jan. 25
Fresno, Cal.	Nov. 18	Apr. 14	Dec. 15	Mar. 4
San Luis Obispo, Cal.	Nov. 19	Apr. 8	Dec. 17	Mar. 17
Los Angeles, Cal.	None.	---do---	None.	Apr. 8
San Diego, Cal.	None.	None.	None.	None.

PROTECTION FROM FROST.

A discussion of methods of protection from frost calls for a consideration in great detail, not only of the several approved processes of protection and the extent to which they can be applied in connection with the various crops, but also the value of these processes under the varying conditions of local topography, local climate, and local soil conditions.

It is evident, in the light of the foregoing statements, that fruit and vegetable growers in rolling and hilly country, or when located in the lee, as regards west to northwest winds, of bodies of water, can, to some extent, so place in their earlier and tenderer fruits and crops that they will, in the first instance, avoid the lowlands and valleys, and in the second take advantage of the more moderate and moist air that crosses water surfaces. For further protection the grower must depend upon artificial appliances. These appliances are designed to produce the following effects or results: To prevent a rapid radiation of heat from the earth; to charge the air with moisture; to warm the air; to create artificial drafts, whereby the air is mixed and the cold air is not allowed to settle to the surface of the earth; or to actually cover or roof in plants.

DEVICES FOR PREVENTING RAPID RADIATION OF HEAT.

Devices designed to prevent a rapid radiation of heat from the earth include screens which can be drawn over plants, vineyards, and groves; loose substances with which low plants may be covered; and smudge fires built to the windward of areas for which protection is desired. These devices, which of a necessity are operative for very limited areas, are described by Professor Hammon in Bulletin No. 23, Weather Bureau, as follows:

Glass screens.—In greenhouses and hotbeds advantage is taken of the peculiar property of glass which allows the heat rays of the sun to pass through it, and is almost impervious to the dark *heat* rays from the earth and plants. This is one of the most perfect screens possible, since it not only prevents the loss of heat by radiation, but receives and retains the heat from the sun. The expense precludes its adoption except for the protection of valuable plants and flowers.

Screens of other solid materials have been quite extensively used in protecting vineyards and citrus groves where intense cultivation is practiced and where the location of the groves near an excellent market admits of profit even with expensive methods of cultivation.

Cloth screens.—In Italy and portions of France screens made of muslin strung on wires stretched on poles above the tops of trees or vines have been used extensively. These screens are drawn on nights when frosts are probable, and are pushed back during the day. When the season has advanced so far as to preclude further danger, they are taken down and stored. Of course such a plan could be operated only on a very limited scale, and would then be expensive. This plan has been recently successfully tried in the orange groves of southern California.

Lath screens.—During the past few years screens made of laths fastened to ordinary telephone wire (the spaces between them being about the width of the laths) have been extensively used in Florida. These are spread over a frame erected above the trees or plants. The screens serve not only as a fair protection from frost, but also as a shade from the hot sun. When no longer needed, they can be rolled up and stored away for preservation. At first thought it would seem improbable that a screen covering only half the space (the spaces being as wide as the laths) would afford much protection; but when it is considered that laths have considerable thickness it is plain that while only one-half the vertical rays are screened those inclined between the vertical and horizontal are partially intercepted by the edges as well as the faces of the laths. As a matter of fact, about three-fourths of the sky is screened by this means. By placing the laths in north and south directions the direct rays of the morning sun are completely cut off from the orchard, which admits of the temperature rising slowly. This reduces the liability of injury to plants.

Other methods.—Strawberries and other low plants are frequently protected by covering them with straw or other loose substances.

Frequently young potato plants are saved by plowing a furrow alongside and allowing the dirt to bury them.

Cranberry growers in the marshes of Wisconsin flood the marshes when frost is expected. In this case the protection is probably due, for the most part, to the high specific heat of the water, as only portions of this land are submerged.

Smudge fires.—Since on cloudy nights radiation is so reduced as to prevent the formation of frost, many have thought that an artificial obscuration of the sky

by means of dense smoke would be an excellent means of protection. The efforts of this character which have been made have resulted in decidedly varying success. In the wheat fields of the Dakotas excellent protection was obtained in limited areas, while the experience of orchardists in Florida and southern California has not shown such uniform success.

Since it was supposed that the protection resulted from the obscuration of the sky by means of smoke, the best protection was expected from the use of that fuel which would produce the greatest smoke.

In the Dakotas the best and most convenient material at hand was the straw of the previous year's crop, which had been left in the fields all winter and through the rainy spring, until it was quite thoroughly soaked with water.

In southern California and Florida straw was scarce, and where it could be obtained it was much drier than that used in the Dakotas. Consequently tar, crude petroleum, and other similar smudge materials were substituted; but the results have not been, as a rule, satisfactory, although the smoke was equally dense. However, quite successful results were obtained by Mr. Buck, Mr. La Rue, and others in the Vacaville and Sonoma sections by burning damp stable manure in sacks scattered throughout the orchard.

Devices for preventing a rapid radiation at night of the heat absorbed by the earth during the day can be utilized only for comparatively small areas. The screen devices, also, by reason of their cost, are available only for the smaller orchards and vineyards whose products command early market prices; the same may be said of coverings of straw and other materials for strawberries and other low plants. Smudge fires can be used to advantage for orchards, vineyards, and ground plants, and even for the smaller grain fields, and would be particularly efficacious in protecting crops and plants in low or bottom lands over which, on still nights, the smoke from smudge fires would settle. There is no device at present known that will, by preventing a rapid radiation of heat, afford any material protection to the great grain fields of the West and Northwest.

DEVICES FOR ADDING MOISTURE TO THE AIR.

Smudge fires made of damp material have been more successfully used for the purpose of affording protection from frost than those made of tar, crude petroleum, and other dry materials, for the reason that in addition to creating a dense smoke they add to the atmosphere a considerable amount of evaporated water, which, though invisible, serves to retard the radiation of heat. When any considerable amount of water is thus converted into vapor and is distributed in the air which covers the area to be protected, a portion of it is likely to be condensed by the surrounding cooler air and appear in the form of mist, which acts as an agent to prevent the escape of heat from the earth; and the act of condensation sets free in the air some of the heat that has been expanded in the process of evaporating the water contained in the smudge-fire materials. Thus the process is

well ordered throughout. In the first place, an undue quantity of heat, whereby undesirable drafts would be created, is not communicated to the air by the fires, and the final result is that in reliberating, by condensation, the water in the air the particles of moisture remain suspended in the air, and some of the heat of the fires is returned to the air in a form which, to a certain degree, warms it without causing currents which would be calculated to disturb the smoke cloud, or pall, and the moist air which overlie the ground and upon whose presence the safety of the plants from frost depends.

As smudge fires made from damp materials are one of the recognized devices which have been found to be more or less effective in affording protection from frost, the following description of methods which have been adopted for utilizing them in orchards is quoted from Bulletin No. 23, already referred to:

Fires of damp straw and stable manure.—Have the fuel, in small piles, distributed throughout the orchard in advance; the more numerous the piles the better. With the same amount of fuel the best protection is obtained from small and frequent fires, since with small fires the upward draft is reduced to a minimum, and the more frequent the fires the more uniform will be the distribution of heat.

Sacks of manure.—A decidedly preferable method is to pack damp stable manure in common grain or burlap sacks, by which it can be conveniently handled. They should be distributed through the orchards in rows about 100 feet apart and about 50 feet between sacks in each row. When it is found necessary to protect, a small amount of coal oil is poured upon each sack and ignited. It is usually unnecessary to fire more than every second or fourth sack, the remainder being left for later occasions. These sacks will burn with a smoldering fire for several hours.

The amount of heat which is set free by burning one sack of manure weighing about 50 pounds and condensing the water vapor near the earth would be sufficient to raise the temperature 20° in a space 75 feet square and 25 feet deep. If one-fourth of this heat remained within this region needing protection, which seems to be a reasonable estimate, ample protection would be obtained for almost any ordinary conditions.

Bales of wet straw.—Mr. T. A. Morrison, of Riverside, Cal., suggested the use of a similar plan, in which bales of wet straw were substituted for manure. This plan has been tried with fair success. One hundred pound bales are cut in four pieces, a tie wire being left about each piece, and if properly dampened will burn with but little care, causing a small smoldering fire.

Prunings.—The prunings of the trees, which are usually removed shortly before the period when frosts are likely to do their greatest injury, are excellent smudge material, and should always be preserved for this use. They should be piled in open spaces throughout the orchard or vineyard and burned at times when protection may be needed. The best results will be obtained from as small fires as will result in burning the prunings.

Portable smudge fires.—A number of excellent devices have been tried, in which the fires were built upon some vehicle by which they could be moved about the orchard. The advantages of this plan are several.

First. The fire can be moved to the section where most needed, which is generally along the windward side of the orchard.

Second. The loss of heat by an upward draft is almost entirely prevented, since the fire does not remain in one position long enough to establish such a draft. On this account much larger and, consequently, fewer fires, with equal efficiency, are possible.

Third. There is a much more uniform distribution of heat and smoke throughout the orchard.

As smudge fires prepared and applied in the manner above indicated are inexpensive, to a degree effective, and easy of operation, they constitute the method which for practical use and utility seems best adapted for general use in orchards, vineyards, gardens, truck farms, and small fields, as a protection from frost. Among the many devices that have been employed for holding and carrying these fires the following has been used with especially satisfactory results by a fruit company of Visalia, Cal., and will doubtless prove equally effective in other sections:

Wire frames (chicken-yard fencing) are built on low truck wagons and stretched from four wagon stakes and heaped over with wet manure. Dirt is then thrown on the wagon beds to protect them, and pots of burning tar are set underneath the straw roof. A barrel of water on the wagon is used to keep the straw wet. The wagons are driven about and do the best work, as they can go wherever the most needed. The smoke and vapor are carried to the rear as the wagon moves, and, being carried at once out of the rising heat, fall close to the ground in a long, white trail. At daylight 400 acres of orchard are covered with a white fog, extending from the ground to about 20 feet high. Similar fires are used as stationary smudges, the wire netting being stretched between four stakes driven in the ground.

Crude petroleum, supplied to a burner from a tank on a wagon or truck, has been used with success by Mr. R. H. Howard, of Riverside, Cal. Over the burner is placed a large wire basket containing mineral wool, which is kept saturated with water supplied by a tank, also placed upon the wagon. Wet straw and manure can also be used. The apparatus can be driven where most needed. The burning coal oil gives off a dense smoke, while the wet mineral wool furnishes a vast surface from which evaporation can take place, and at the same time the material will not be consumed.

DEVICES FOR CHARGING THE AIR WITH MOISTURE.

Aside from the smudge fires, Bulletin No. 23 quotes devices which have been adopted to charge the air directly with moisture, some of which are constructed on a plan too elaborate and expensive for general use.

The first operation involved is the evaporation of water in large quantities, either by means of evaporating pans or by the use of boil-

ers with connecting pipes whereby steam can be generated, carried, and thrown into the air in different parts of a field or orchard.

Spraying and sprinkling plants in times of threatened frost has also been attended with some degree of success. At the Everest ranch, at Riverside, Cal., sprinklers are placed at the top of 50-foot masts, which fill the air with a very fine spray. In nearly every instance the protection against frost by this method has been sufficient.

Sprinkling gardens before sunrise on frosty mornings has proved of decided value.

Secondly, irrigation, which of necessity can be used only to a very limited extent and in certain favored localities, has been found to afford very complete protection from frost in fields, orchards, and vineyards which are equipped with irrigation ditches, and in the cranberry districts of Wisconsin perfect protection from damage by frost is secured by flooding the cranberry marshes the day before heavy frost occurs in that section.

In no case should water be allowed to stand in orchards, vineyards, or gardens during the early growing period while danger from frost is possible, for the reason that water unduly promotes the growth of young plants and thereby renders them more susceptible to damage by frost and cold than are the plants whose growth is retarded.

The third operation involved in the California experiments was a direct heating of the air by means of fires. It was found that large fires caused a strong upward draft of air, which had the effect of drawing in cooler air from the sides, and that well distributed small fires were attended by very satisfactory results.

The following final report of the Riverside Horticultural Club covers the results of experiments made during the winter of 1897–98:

With the assistance of fifteen or twenty citizens interested in the study of the points involved, a most complete test has been made of the many different methods employed to prevent frost damage. With such a force of competent and impartial observers it was possible to secure data of much value in forming an estimate of the efficiency of the various plans made use of. Careful comparison was made between those orchards where no work was done, and where no direct effect of the fire was probable, and those where the different methods were being tried. As indicated by our partial report at the last meeting of the club, these tests were in some particulars eminently satisfactory as showing the way to definite conclusions.

The exceptionally long period of cold following gave additional opportunity to verify the first conclusions reached, and subsequent investigations made by ourselves, as well as by other citizens who have awakened to the possibility of protecting their property, strengthened and confirmed the opinion formed as the result of the tests already partially reported upon.

Some theories are proved to have little practical value, and members of your committee have modified their views somewhat in consequence. No preconceived notions have been allowed to stand in the way of a thoroughly practical study of the facts as they exist, to the end that the growers may not, for the

lack of definite knowledge as to the directions their efforts should take, neglect reasonable precautions hereafter to insure the safety of their crops. These, however, are our conclusions:

First. There is no doubt whatever that the temperature of our orchards may be materially raised by the use of dry heat.

Second. The radiation of the earth's heat can be considerably lessened by moist smudges, when these are started early enough and are properly managed.

Third. The possibility of raising the dew-point on one of the dry, cold nights peculiar to our climate sufficiently to prevent damage, by means of steam-producing apparatus, seems impracticable.

Fourth. Fruit and trees can undoubtedly be saved, even in the coldest sections, by covering them with cloth or matting, but the expense involved makes this method impossible on the part of the ordinary grower.

Fifth. It is found that the temperature in an old seedling grove, or where tall wind-breaks afford to smaller fruit trees a like protection, the temperature is almost invariably 1° to 2° higher than in exposed orchards in the immediate neighborhood. This fact seems to thoroughly upset the theory, strongly held by many intelligent growers, that the tall, well located wind-break is a disadvantage, the contrary seeming to be the truth.

Sixth. It is found that the temperature 20 feet above the ground is 1° to 2° higher than at the surface, and that, as a rule, when the cold is severe enough to injure the ripest fruit 50 feet from the ground there is almost invariably a temperature above the freezing point of water.

Professor Zumbro, who has given especial attention to this matter, finds that at the height of 50 feet the temperature is from 5° to 10° higher than at the surface when the air is not in motion. When there is any considerable breeze it varies but little.

Seventh. Our conclusion is that, all things considered, the coal baskets, sufficiently numerous, will prove the most satisfactory and effective means of warming the orchards yet made use of. It is true the oil pots make a far hotter fire, and are neither expensive nor difficult to manage; but the deposit of lamp black upon tree and fruit resulting from their use condemns this system for general use.

As to the value of smudging the members of your committee are not so well agreed. Because of less sharply defined results, we find it more difficult to come to definite, at least uniform, conclusions. But, under certain conditions, we are convinced that, properly used, it may be made a valuable means of protection. We think this especially true in localities where the temperature falls only a little below the danger point, and where there are considerable solid areas of young orchards exposed. Here it will work well if the protection is made general. But where the danger is considerable we think it wise to be prepared to use dry heat, even where in connection with the smudge. The benefit from smudging is probably as much from its protecting fruit and trees from the sudden rays of the morning sun after a freezing night as from modifying temperature during the time of danger.

Experience demonstrates that flooding or running water in connection with dry heat or smudging is a valuable adjunct. One of the committee who has been testing this matter carefully for three years is disposed to think that the direct benefit from running water is overestimated by a majority of growers. Its value in putting orchards in condition to withstand quite severe weather safely is questioned, but the committee are inclined to think that entire dependence upon this method will occasionally result in serious loss to those who trust to this means alone, especially when used in young orchards.

As to the number of baskets needed when coal is used, we find that the most decided and satisfactory results have been gained where from twenty to fifty coal fires have been used to each acre. If intelligently and energetically used this plan will never fail, except when the mercury drops below 24° for a long while, and even then it is believed the larger portion of the crop may be saved if anything like a general use of such fires be secured. The smaller number of fires named has, in numerous instances, and even when a man was working alone, secured a rise of from 3° to 5°, and saved a crop. Can it be doubted that 50 fires per acre, used in every orchard, would save both trees and crop on the coldest night ever known in California's history?

To equip an orchard with 50 baskets to the acre means an outlay of only a little over $5. The fuel to run them one night costs from $2.50 to $3. If a crop of navels upon it is worth $400 it will pay well to spend in labor and fuel $4 per night, or 1 per cent on the value of the crop to insure its safety. In the orange region of southern California it is not unusual to have more than two or three nights in a season when the fruit is in danger. But even if, as in the present season, the period of cold is more extended will it not pay to expend at least as much as one pays for his irrigating water to insure the safe maturing of a crop it has cost him a year's labor and heavy expense to produce? The conclusion is obvious that we have only to provide for the insurance of this sort of property exactly as we would in the case of that liable to destruction by fire, to be enabled to follow the business of orange and lemon growing with the certainty of having perfect fruit to market at the season's end.

While the practicability of protecting our orchards from frost seems established, the problem of the most economical and scientific means of accomplishing this is probably yet to be solved. However well the wire baskets may serve us now, there will doubtless be improved method for burning coal, and even other material may be found that will serve the purpose better; and while wet straw seems to be at present the most available for smudges, doubtless, when the need is made known, chemists will find some vapor-producing material more compact, efficient, and economical. Hence we recommend that the club appoint a permanent committee to continue these investigations.

GENERAL OBSERVATIONS AND PRACTICAL RESULTS.

WORK IN TEXAS.

In a paper read before the annual meeting of the Texas Farmers' Congress and Texas State Horticultural Society in joint session at College Station, Tex., July 26, 1899, Mr. Joseph L. Cline, observer, Weather Bureau, submitted the following remarks on "The use of frost and temperature warnings in protecting fruits and truck gardens:"

It has been proved that a good way for protecting berry crops is to cover with hay. Berries are improved by mulching. Prairie hay used for this purpose can be cut and placed in the fields, ready for covering the plants when frost is indicated, at a cost of about one dollar a ton. Ten tons per acre are considered sufficient for protection against any freeze. The cost of labor for covering the plants would be about two dollars an acre additional. As the hay is required for mulching, the cost is considerably lessened when the hay is utilized for that purpose. Berry crops which were covered prior to the freeze of last February yielded four times as much as unprotected plants, and unprotected plants were generally killed. Cabbage, potatoes, and other vegetables

can be protected from frost by hay, and the increased yield will more than pay for the expense involved. Rice growers along the east Texas coast can raise vegetables throughout the winter on their rice plantations by protecting them from frost. The plantations can easily be flooded during freezing weather, which is always of short duration, and the water would protect and do very little or no damage to the vegetables, which are remunerative during the winter season.

* * * * * * *

The method of burning stakes soaked in petroleum or kerosene oil throughout the fields can be applied not only to the protection of maize, but fruit orchards as well. By means of burning stakes fields will be enshrouded, when the wind is light, in a continuous cloud of smoke. This will, in many instances, prevent frost from occurring. The stakes must not be set so close to the fruit trees that the heat will affect the branches.

* * * * * * *

In France frost bells are sounded when frost is expected, and grape growers protect their crops by pouring tar on the ground among the vines and setting it on fire. The smoke thus made often saves the crop of an entire vineyard. Bituminous coal burned in baskets is less expensive and has been successfully used. Forty to fifty baskets of coal per acre are required, and the more used the better would be the result. Strawberry crops can be protected in this manner at a cost of $4 to $6 an acre.

* * * * * * *

The following letter from Capt. B. F. Johnson, president Gulf Coast Horticultural Association, gives an idea as to what can be done in protecting crops [by showing what was done in February, 1899]:

"I was in Galveston, Tex., on the morning of February 11, 1899, and upon hearing that a freeze was expected I called at the United States Weather Bureau office, and was informed that the hardest freeze of the season was threatened. I immediately gave orders over the long-distance telephone to my employees at Arcadia to cover up all hotbeds and also all plants already transplanted. All the plants in hotbeds were carried through the freeze by protection and saved from injury. I had about 7,500 cabbage plants set out, and on account of my employees not going to work as directed we did not reach these until ice had formed over some of the plants. I took the first train out of Galveston, which was in the afternoon, and upon my arrival at Arcadia we continued protection, notwithstanding the fact that ice had already formed over the plants. We placed a handful of hay over the north side of about 1,000 cabbage, and at least 750 of these came through the freeze uninjured, while all the unprotected were killed. The varieties protected were the Jersey-Wakefield and Henderson-Succession, and I could see no difference between the two after the freeze. Had my employees carried out the instructions given them in the morning I would have saved 75 per cent of the 6,500 unprotected plants which were killed, and this would have yielded me about $150 net. I had 1¾ acres of Irish potatoes planted in January, which were up to fair stand. These were not protected and were all killed. I had about one-sixteenth acre planted with hand planters the 1st of February. These were protected with vines from old sweet potatoes grown on the ground the previous year. Fully one-eighth of these came up after the freeze with this light protection, and thus enabled me to put the first potatoes on the Texas market, and they brought me $3 a bushel net. If I had been prepared and protected the 1¾ acres planted in January I would have received a nice little sum, and more than ten times the

cost of protection. Strawberries not mulched were virtually killed, while those mulched made an average crop. The great value of protection to berries in case of freezing weather has been forcibly illustrated. The entire crop of early berries (the only berry crop which pays well) would have been a total failure without protection."

* * * * * * *

The following communication of Mr. M. M. Cox, president of the League City Truck Growers' and Farmers' Association, League City, Tex., gives further the manner and value of protecting crops from the frost and freeze during the abnormally cold weather of February 12 and 13, 1899. Mr. Cox says:

"I had two acres of strawberries—Noonan and Mitchell varieties—which were blooming and fruiting nicely. In anticipation of a freeze we placed hay for mulching purposes between the rows, which were about 3 feet apart. Upon the receipt of the telegram, Saturday morning, February 11, 1899, announcing the approach of a freeze, I got word to the man who had charge of my berry crop. He immediately went to covering the vines with hay previously placed in the berry patch and succeeded in covering 20 rows. The vines of the protected berries were practically uninjured, while the unprotected plants were frozen to the ground and many of them totally killed. From these 20 rows, about one-fourth acre of protected plants, we picked berries nearly three weeks before the unprotected berries were ready for market. The one-fourth acre of protected berries yielded me more than the remaining 1¾ acres altogether. In fact, we got the bulk of our crop—a little more than $300 net—off this one-fourth acre. If we had used more hay for covering we could have saved the blooms and berries already on the vines notwithstanding the severity of the freeze, as some berries were saved as it was.

"Others in League City protected berry crops with the same results as those which attended our protection.

"Hotbeds protected for ordinary freezes were killed, but some put extra covering on the beds, using blankets, and thus saved their plants without injury. Those who protected in this manner were setting their plants in the field while others were sowing seeds in the hotbeds, and were thus at least four weeks ahead with their crops for the market.

"The warnings of the Weather Service promise much for the coast country."

Mrs. C. W. Benson, Alvin, Tex., in writing of protection of crops, says:

"In past seasons the advance notices of frost and freezing weather given out by the United States Weather Bureau have been distributed among the truck growers by messenger service, and by this means have saved our planters who protect their crops thousands of dollars.

"In 1899 all crops were carried through safe until the severe freeze of February 11, and in several instances tender crops were successfully protected and carried through this freeze even when the mercury stood at 4 degrees above zero or 28 degrees below the freezing point. The principal crops protected were tomatoes, cucumbers, cabbage, and strawberries. The three first mentioned were still in hotbeds or cold frames, and were protected by covering with hay to a depth of about 1 foot, over glass or canvas. Berries were protected by turning over mulch and covering up plants from 4 to 6 inches in depth. This mulch had already been placed between the rows to retain moisture and keep the berries clean on plants. This protection did not save blooms, but the plants so protected came through the freeze with much less damage and produced more fruit than the unprotected plants. I judge that the protection was fully worth $30 per acre, as prices of berries remained firm all through the season on account of general scarcity of fruit. About 25 acres were thus protected from the freeze.

"The value of the tomato crop saved through protection from this freeze averaged fully $125 per acre in spite of the subsequent drought, which extended from February 14 until after the crop was marketed, with the exception of a few slight showers, which were insufficient for general growth of plants. With rain as in an average season the crop would have paid $250 per acre, and enough plants were carried through this freeze by protection to set 17 acres.

"About 200,000 cabbage plants were carried through this severe freeze of February and successfully transplanted. These would have netted the growers $8,000 with sufficient rain during the growing period. The value of the cucumber crop from actual results was about $1,000, which would have increased 1,000 per cent in an ordinary season, but they were held back for lack of moisture until the north Texas crop was on the market, as cucumbers which sold at from 40 to 50 cents per one-third bushel box would have netted from 75 cents to $1.25 per dozen.

"Protection of crops enables the grower to generally reach the first markets, but without the advance notices of frost and freezing weather it would be impracticable to protect; our people would bear the expense of the warnings rather than do without them."

WORK IN LOUISIANA.

Probably the most extensive experiments for the protection of citrus fruits in the Southern States have been conducted in a large orange orchard near Diamond, La., under the management of Mr. Hewitt Chapman. In the Report of the Louisiana Section of the Climate and Crop Service of the Weather Bureau for February, 1899, this grove and some of the methods employed to protect its crop are described by Mr. Alexander G. McAdie, of the Weather Bureau, as follows:

The orange orchard is about 43 miles south of New Orleans, near the river, and about 20 miles north of what is considered the safe orange belt of Louisiana. The grove covers 130 acres and contains 15,000 trees, including Creole Sweets, Mandarins, and Satsumas; sweets budded on Trifoliata and Pomeloes on sour stock. These young trees withstood the frosts of January 1, December 4, 9, 10, 11, 1898. The writer visited this grove on December 11, when ice one-quarter inch in thickness could be found along the roads and in exposed places. The orchard is irrigated by means of river water carried over the levee by a direct-acting pump, 6 by 8, worked by a 40-horsepower boiler. The pump dispenses with belt, pulley, and engine. To flood the whole orchard of 50 acres to a depth of 1 inch would require 5,650 tons of water, or 1,357,700 gallons. Barely one-hundredth part of this quantity of water is required. The water pumped from the river over the levee goes into a main ditch, from which run subditches, each subditch feeding two small inlets through the orchard between each line of trees. The distance between the trees averages 12 feet. The middle ground is cultivated and level so that the water can run in a steady stream to and out through the quarter drain into the drainage ditches which are 1 acre apart, running parallel into the rear canal.

The grove has a northern exposure; fronts on the river; the land runs west and east, falling slightly from the river; and there are no wind breaks. It is also proposed at this orchard to experiment with the various types of smudges. One portion, about 10 acres in extent, will be protected by smudge fires, some portable and some stationary. In another section, covering about 30 acres,

the bodies and lower portions of the orange trees are to be wrapped with native hay. In the third section the dirt is to be hilled up about the trees; while in the fourth the trees are to be wrapped with floor matting.

In the methods described above for the protection of trees it was thought that colder weather than that of February, 1895, was not likely to occur again. It appears that the pumping plant, on which great reliance was placed, did successfully hold back the trees up to about February 2, 1899, but the extremely warm weather of the 3d, 4th, and 5th started the sap and forced growth. It is doubtful, however, whether water protection is adequate under such severe tests. All the orange trees above ground appear to have suffered badly, in nearly every case the tree being killed. Those trees which were wrapped with matting likewise suffered and the protection again appears to have been inadequate to withstand such extreme cold. In the section where the earth was hilled up around the trees the results were very satisfactory. It is thought that about 4,000 trees have been saved by this means. An inspection of the orchards as late as March 5 indicates that notwithstanding the severity of the test, there is reason to believe that orange trees can be protected successfully by hilling up the earth.

The devices employed in Louisiana, as described above, were found to be an effective protection against damage by frost, and the protection afforded by irrigation and by wrapping the bodies and lower portions of trees with hay was sufficient for ordinary freezes. During the unprecedented cold of February, 1899, the section in which dirt was hilled up about the trees was the only one in which trees escaped serious injury.

WORK IN FLORIDA.

Devices used in Florida for the protection of fruit, vegetables, and berries from frost and light freezes conform in character to those tested in the California experiments, in which fires were found to be the most effective means of protection, and appliances for adding moisture to the air were successful only to a slight degree.

As artificial appliances are totally inadequate to add to the atmosphere any very appreciable amount of moisture, it is evident that methods which may have been found ineffective in the dry climate of California would possess value in localities where their office is confined to adding moisture to an already moist atmosphere. The Gulf and South Atlantic Coast States, and in fact the country generally from the Mississippi Valley to the Atlantic seaboard, possesses a moist atmosphere, and the fruits and tender vegetables of these districts can, therefore, be the more readily protected from frost by devices which add moisture to the air. As a means of adding moisture to the air, irrigation should be more effective than in California; and in localities where this method can be used protection will be assured except against hard freezes.

Owing to the comparatively inexpensive character of the materials used in damp smudge fires, they seem the best adapted for common

use in groves and gardens. Berries and other low plants can be protected with but little expense by coverings of straw and other light materials. In Florida pine needles are successfully used to protect strawberries. Devices for actually heating the free and open air are expensive, and their value is dependent solely upon a comparatively still air and small, numerous, and well-distributed fires.

In Florida many experiments have been made with a view to adopting devices which will protect citrus trees and fruits from injury during the periods of severe cold which at times visit that section. The fact that these periods of cold are infrequent does not relieve the grower from the necessity of providing for their occurrence. Effective measures of protection necessarily vary with the conditions that confront the grower. Protection can be made effective by tents, coverings of cloth, sheds, and one-half and one-third lath shelters and their modifications, that retain heat from open wood fires and sheet-iron stoves. The expense of these methods is obviously too great to make their employment, except for choice varieties of fruit, commercially profitable.

In this State protection of citrus trees and fruit in the open field is usually afforded by what is known as "firing." This method is effective, except in the presence of severe freezes, provided rain or strong winds do not accompany the cold. Heavy rain will, of course, extinguish the fires, and strong winds will not permit a necessary accumulation of warm air about the trees. When no rain is falling and the wind is light, fires must be built on the windward side of the trees, so that the heat will be carried through the trees, and the number of the fires must be regulated by the degree of cold. In this form of protection the grower must lay in and place his materials for "firing" before the winter season sets in, and must, during the season, be on guard to utilize the materials (usually wood) that he has accumulated. When local indications point to dangerously low temperatures, or when warnings have been received from the Weather Bureau, he should watch his thermometers, exposed in the groves, and be prepared to "fire" with a rush when the mercury sinks to the danger point, 28°, with a downward tendency. There is no time on the eve of a freeze to collect firing materials, and the grower must not desert his grove for bed or board so long as danger by cold continues. He will have his hands full keeping his fires going, even if he has sufficient material for that purpose. Material should be available for at least three nights. Cold spells seldom last longer. The danger increases with the length of the cold spell. Trees and fruit become chilled and the last night of a cold spell is likely to be the one that does the damage. The freeze of December 23–27, 1906, lasted five nights, and in some sections the last night produced the

lowest temperatures of the period. In some localities growers had exhausted their supply of firing material and, as a consequence, suffered a loss of fruit and young trees.

In cases of severe cold fruit can not be saved, and trees can be preserved only by banking up the trunks by sand or soil. Small trees can be buried, and older trees can at least be protected above the parts where they have been budded.

The damage that will result from a given degree of cold varies with the condition and size of the tree. Well-grown trees in a dormant condition will stand without serious injury, except to foliage, a temperature of 18° to 20°, provided it is not too long continued. When the sap is flowing these temperatures are likely to kill growing trees, and are apt to cut small, unprotected trees to the ground. Well-grown trees in a dormant condition will stand without injury a temperature of 26° of continued cold, and can for several hours stand temperatures of 22° to 24°. When the sap is flowing these temperatures continued for several hours will be more or less injurious. Cold that will freeze fruit will injure small trees. In the freeze of December 23–27, 1906, temperatures in parts of the orange belt fell to 20°, and even to 18° and below on one and in some cases two nights without injury to large trees, except to foliage, and with a surprisingly small loss of fruit and small trees. This freeze had been preceded by about two months of rainless weather, and found the trees in a dormant condition. The destructive freezes of preceding years usually followed periods of wet weather that promoted a flow of sap.

Slight protection can be afforded groves by running water in irrigation ditches, or by spraying the trees with water, each process imparting some warmth to the air that is derived from water that is much warmer than the air during a cold spell. As it is not, as a rule, possible to run water in ditches made in Florida sand the irrigation method possesses little value in that State. There are precautionary measures that may in some instances be employed to advantage. In setting out a grove there should be left, where possible, forest trees on its north and west sides where they will act as windbreaks during cold spells. High fences are sometimes employed as windbreaks. Theoretically cold air drains from high to low ground, and groves on elevated sites are more safely located than those on low ground. Protection should also, where possible, be given by water surfaces on the west and north of groves.

FROST SEASONS FOR SEVERAL CROPS.

Damaging frost is likely to occur in northern Florida as early as the middle of October and as late as the early part of April. During the winter period severe freezes occasionally occur in middle and

northern portions of the State, and at rare intervals freezing temperature is experienced to the southern end of the peninsula. Except on the keys and near the coasts of the peninsula south of about the 27th parallel, pineapples can not be safely grown in the open field. When grown in the interior north of that parallel, or on the more northern coasts, they should be afforded means of protection. This is usually furnished in the form of one-half or one-third lath shelters built over the field. Fancy varieties are often grown under glass. Strawberries are subject to damage during the winter and early spring; oranges from December to the early part of February, and orange trees are subject to damage by freezes during December, January, and February. Garden truck is especially subject to injury from January 1 to April 15. In the trucking districts of the interior of the Gulf and South Atlantic States the period during which damaging frost is likely to occur extends from October to April, inclusive.

Temperature conditions are of great importance in the culture of rice, and a severe frost late in October may injure the grain so much as to cause its discoloration. Owing to the manner of growth and culture of this crop and the character and extent of the fields it covers, practically nothing can be done to avert injury from heavy frosts that may occur about the time of harvest.

Tobacco growers may protect their plant beds by coverings, but during the critical periods following transplanting and approaching harvest severe frost will ruin plants in fields which are too large for the successful application of the devices herein described. During the maturing season tobacco plants in large fields can be given protection from damage by frost only by cutting and "frost heaping" on the eve of the occurrence of frost.

Cotton will be seriously injured by low temperature early in the spring whether frost occurs or not. Owing to the area covered by the fields, no adequate means of protection against cold and frost, suitable for general use, has been discovered.

In Louisiana sugar cane is planted in the fall, winter, and early spring. The plants are often killed or seriously injured by frosts in the spring and in October and November. When frost is expected, the cane can be cut close to the ground and "mattressed," the product of three rows being so laid together that the leaves of one armful will cover the butts of the preceding; being thus thatched the canes are protected from frost and will keep in this state for several weeks without injury.

In the cranberry districts of New Jersey, Cape Cod, and Wisconsin, and in limited areas in other parts of southeastern New England, and the upper lake districts, which are devoted to cranberry raising, flooding is the only effective means of protection against damage by frost. This crop is confined to swamp or bog lands, and when danger from

frost is apprehended appliances for letting in a sufficient quantity of water to keep comparatively warm and moist the lower stratum of air are usually provided.

Young potato and tomato plants can be saved from damage by frost by a covering of dirt. In large patches this can be accomplished by plowing a furrow alongside the rows of plants and allowing the dirt to bury them.

Gardens can be protected from frost by coverings of straw, paper, or other loose and light materials, or by damp smudges; and the latter device appears to be, in point of effectiveness and cost, the best adapted for general use not only for small areas devoted to gardens, but also for orchards, vineyards, and fields. The effectiveness of any method of protection is of course decreased as the area for which protection is desired is increased, and there is no process yet devised that can give adequate protection against frost and cold in the great grain fields of the country.

CONCLUSIONS.

Experiments and observation seem to establish the following facts: The danger of damage from frost can be materially lessened by placing early and tender plants on high grounds and crests, and hardier plants in low grounds and hollows. When ground can be selected in the lee, or to the south and east of considerable bodies of water, the danger will be further lessened.

In the dry climate of the citrus fruit region of California and in the promising fruit districts of Arizona small and numerous fires, preferably of coal burned in iron baskets, have been found to be the most effective device used for protection against frost; second in point of utility may be placed irrigation; and the practicable process which affords the least protection in that district appears to have been smudge fires. As before stated, the unsatisfactory character of the experiments made with smudge fires may be attributed to the fact that artificial devices are inadequate to charge dry air with sufficient moisture to prevent loss of heat from the earth by radiation. Reports of California experiments do not distinguish between frosts and freezes. Owing, however, to the usually moderate character of the cold periods of that section the devices referred to would doubtless be equally effective against frosts and freezes.

In the orange-growing districts of the South irrigation affords the most effective protection against frost, while in sections where this process can not be employed damp smudge fires for frosts and light freezes, and wood fires for moderate freezes, properly handled are best adapted to general use. Severe freezes which kill or injure the trees can be guarded against only by housing the trees and heating

the confined air, or by "banking up" the trees with dirt or sand. The latter method is comparatively inexpensive and is therefore the better adapted to general use.

In all sections loose coverings may be used to advantage for protecting low plants, and damp smudge fires may be profitably employed in gardens, orchards, and small fields.

It is important to remark in this connection that none of the artificial devices described should be employed during the growing season except on the immediate eve of threatened frost. The necessity for observing this precaution becomes apparent when the fact is considered that any appliance for protecting plants from cold will maintain abnormally high temperatures when average weather conditions prevail. The effect of this process would be to force the growth of the plants and render them the more susceptible to damage by the late frosts of spring. In all sections and localities reached by the forecasts and special warnings of the Weather Bureau action to put into operation protective measures can be safely deferred until warning notices are issued. The Bureau has made provision for a distribution of frost warnings which is limited only by the resources of the telegraph, telephone, mail, train, and signal services, and agriculturists whose location renders possible the receipt of the warnings by any of the means referred to should adopt measures for securing them. That action in this direction would be profitable is shown by the testimony of those who have received substantial benefits from the warnings.

104

O

Reserve

Issued December 14, 1910.

U. S. DEPARTMENT OF AGRICULTURE.

FARMERS' BULLETIN 104.

Rev.ed.
follows

NOTES ON FROST.

BY

E. B. GARRIOTT,
Professor of Meteorology, Weather Bureau.

Revised July, 1910,

By ALEXANDER G. McADIE,
Professor of Meteorology, Weather Bureau.

JAN 3-1910
U. S. Department of Agriculture

WASHINGTON:
GOVERNMENT PRINTING OFFICE.
1910.

LETTER OF TRANSMITTAL.

U. S. DEPARTMENT OF AGRICULTURE,
OFFICE OF THE WEATHER BUREAU,
Washington, D. C., August 2, 1910.

SIR: I have the honor to recommend that the manuscript herewith be published as a revised edition of Farmers' Bulletin No. 104.

In compliance with directions contained in your letter of March 29, 1899, a paper on the subject of frost, prepared by Prof. E. B. Garriott, Weather Bureau, was submitted and published as Farmers' Bulletin No. 104. This bulletin furnished valuable information regarding methods of protection from frost. Methods advocated in previous publications were mainly the results of experiments conducted in California orchards. Bulletin No. 104 was designed to be of value in all sections of the country rather than in restricted districts.

As the California experiments quoted were perhaps the most thorough ever made in this country, a description of the methods employed and the results obtained is reprinted in brief in this paper, together with results of similar experiments made in the citrus-fruit and truck-growing districts of the Gulf States, where varying climatic conditions have been found to produce somewhat different results. Additional information personally acquired by Professor Garriott during a visit to Florida in the winter of 1906–7, and certain climatic data, also form features of this paper in its revised form.

The development of orchard heaters and other devices now successfully used in saving crops from injury by frost necessitates a new issue of this bulletin. Owing to the death of Professor Garriott, the revision has been undertaken at my direction by Professor McAdie.

Your wishes that the subject be treated in a popular way and divested of all technicalities, and made applicable to all portions of the United States, have been complied with so far as the nature of the subject will permit.

Respectfully,

WILLIS L. MOORE,
Chief, Weather Bureau.

Hon. JAMES WILSON,
Secretary of Agriculture.

CONTENTS.

	Page.
How frost is formed	5
Seasons of frost	7
When to expect frost	10
Influence of soil and vegetation on minimum temperature	14
Dates of early autumn and late spring frosts	15
Protection from frost	17
Devices for preventing rapid radiation of heat	17
Glass screens	18
Cloth screens	18
Lath screens	18
Paper cover	18
Other methods	18
Devices for adding heat to the air	19
Devices for adding moisture to the air	24
Portable smudge fires	25
General observations and practical results	26
Work in Florida	27
Work in Missouri	29
Distribution of forecasts in California	30
Miscellaneous	31
A two years' study of spring frosts at Williamstown, Mass	31

ILLUSTRATIONS.

FIG. 1. Chart showing average date of first killing frost in autumn	8
2. Chart showing average date of last killing frost in spring	9
3. Diagram of Kunkel orchard, Wichita, Kans	22
4. Diagram of plat of Thomas Orchard Company, Wichita, Kans	23

NOTES ON FROST.

HOW FROST IS FORMED.

The atmosphere of the earth always contains more or less moisture in an invisible form. When at a considerable elevation above the earth this moisture, or aqueous vapor, is condensed, clouds are formed; when the process of condensation is more active and the temperature of the air is above freezing, rain falls; and when the temperature of the air is below freezing, snow is produced. When the moisture of the air in immediate contact with the earth is condensed at temperatures above freezing, dew is formed; when at temperatures below freezing, frost is deposited. Frost is, therefore, the moisture of the air condensed at freezing temperatures upon plants and other objects near the surface of the earth.

In the process of frost formation the temperature of the air a few feet above the earth is commonly several degrees above freezing. The surfaces upon which frost is deposited must, however, possess freezing temperatures. The manner in which frost is deposited on plants and other objects is very similar to that in which the air moisture of a room is frozen and deposited upon window glass, the temperature of which has been reduced to freezing by the out-of-doors cold. In the case of the frost on the window glass the process is one that can readily be understood. Some explanation, however, is required of the formation of frost, which requires freezing temperature, when at times a temperature above freezing is registered a few feet above the surfaces upon which frost appears.

There are several processes by which the temperature of plants may be reduced below the temperature of the air which surrounds them. The most important of these processes is radiation, by means of which heat escapes from objects and passes into the surrounding air.

In the frost-forming process heat from the sun which is absorbed by the earth and by plants during the day is lost by radiation at night. During the day the earth both absorbs and reflects the heat received from the rays of the sun, and the lower stratum of air is warmed by this reflected heat. During the night when no direct heat is received from the sun's rays the lower air stratum receives no reflected heat,

and, at the same time, heat which has been absorbed by the earth is radiated, or, in other words, it rises through the overlying air, though not necessarily and entirely as sensible heat; that is, heat perceptible to the senses. In the frost-forming process there is another very important factor in the production of cold; that is, the evaporation of moisture from the earth and from plants. The nature and composition of frost require that the air in immediate contact with the surfaces upon which frost forms shall contain moisture. This is moisture that has been stored in the earth in visible form, i. e., in the form of water, and which has escaped from the earth, not as visible moisture, but by the process of evaporation and in the form of aqueous vapor. Some part of the heat stored by the earth during the day is, therefore, lost in evaporating the moisture contained in plants and in the earth. The heat expended in this process is not sensible, but latent. Heat is rendered latent when it is used to boil away water, and it becomes sensible or liberated heat when the water vapor again condenses into water. It appears, therefore, that the temperature of surfaces upon which frost forms, and of the air in immediate contact with them, is lowered by the evaporation of moisture from the surfaces. The influence of the process does not extend to air a few feet above the ground.

Another method by which plants lose their heat is by convection, by means of which they are chilled by contact with colder air. This process, while important in the presence of freezing air temperatures, can scarcely be considered a factor in the formation of frost proper, which is usually accomplished when temperature observations show the air to be above freezing.

Air arranges itself according to its density or weight. Air in immediate contact with the earth becomes heavier as its temperature is lowered by the radiation of heat and settles in depressions or valleys and over lowlands, causing, at times, frost in low-lying districts, while neighboring higher grounds escape the visitations.

NOTE.—A given quantity of heat absorbed by several substances of different specific heat will produce a different temperature in each; the lower the specific heat of the substance the higher will be its temperature; and besides, different substances have different coefficients of absorption and of reflection. Therefore under the same insolation rocks, clay, vegetation, and other substances come to temperatures that often differ by many degrees.

The air next to earth largely partakes of the temperature of the surface, and in consequence, when the sky is clear, so that incoming and outgoing radiations may progress freely, and the air is but little disturbed by wind, wide variations in the temperature of the thin stratum of lower air may occur over adjacent plots of ground of *precisely* the same elevation but different covering.—Moore's Descriptive Meteorology, page 108.

SEASONS OF FROST.

Generally speaking, agricultural products in some part of the United States are menaced by frost from the time nature begins to extend her mantle of green from the citrus regions of the South toward the grain fields of the North until the verdured area contracts southward before the first winds of winter; from the hour that seeds germinate and buds blossom until crops are gathered; from early spring until late fall. As a rule, the critical periods are during the early growth of plants and when the fruit is in the bloom, and again when crops are maturing. In the first instance vegetation is threatened by the frosts of early spring, and in the latter by the frosts of autumn.

Considering the several agricultural sections in greater detail we find that damaging frost is likely to occur in the north half of the Florida peninsula, the region immediately bordering the coast of the Gulf of Mexico, and in the interior of California from the latter part of October until the early part of April, and in the North Pacific Coast States as early as the middle of October and as late as the last week of April. From December until early in March the frost limit and even the line of freezing temperature may, at long intervals, be extended southward over central and southern Florida. The period of damaging frosts in the interior of the South Atlantic and Gulf States extends from November to April. From October to April the region in which agricultural products are subject to damage by frost is extended to the southern lake region, over the Upper Mississippi and Lower Missouri valleys, Kansas, and Nebraska, and from September to May frost visitations are usually confined to the extreme Upper Mississippi, Upper Missouri, and Red River of the North valleys, and the Rocky Mountain and plateau regions from central New Mexico and the Texas Panhandle northward. (See figs. 1 and 2.)

The occurrence of summer frosts in mountain regions and in extreme northern districts of the country is common. The following are notable instances of summer frosts in agricultural districts:

August 22 to 24, 1890, a frost-bearing cool wave advanced from the Dakotas over the lake region and the interior, or mountain districts, of the middle Atlantic States.

August 28 to 31, 1892, a cool wave advanced from the Rocky Mountain and plateau regions, with frost from New Mexico to the Dakotas, Iowa, Minnesota, northern Wisconsin, and northern Michigan.

August 8, 1904, frost damaged vegetation in low-lying lands in the States of the upper lake region.



FIG. 1.—Chart showing average date of first killing frost in autumn.

104



FIG. 2.—Chart showing average date of last killing frost in spring.

Bulletin P, of the Weather Bureau, gives a chronological statement of the more important cold periods that have been experienced in the United States, and contains charts that present conditions that preceded and attended the principal cold waves and frosts of the fifteen years from 1888 to 1902.

WHEN TO EXPECT FROST.

As the conditions which produce damaging frost are subject to modifications which are as numerous as the kinds of crops raised, and as varied as local topography, local climate, and local soil conditions in the various sections of the United States, the writer in attempting to treat briefly a subject which is so diversified in its aspects is confronted with many difficulties. Suffice it to say that with other atmospheric conditions favorable for its occurrence frost may, as a rule, be expected when temperature, as reported by stations of the Weather Bureau, falls to a point 8° to 10° above the freezing point. As previously stated, the surfaces upon which frost is deposited must be at or below freezing temperature, although the temperature of the air a few feet above may be several degrees above freezing. It is the temperature of the air, in some instances many feet above the ground, that is given by the Weather Bureau observations. Another atmospheric condition favorable for the occurrence of frost is a clear, cloudless, and comparatively calm night. Clouds retard radiation or loss of heat from plants and act as a screen in preventing the heat collected from the sun's rays during the day from escaping into the upper air. When clouds are not present and a withdrawal of the sun's rays causes a rapid cooling of the air at moderate elevations, the warmer air which collects near the surface of the earth during the day rises, and the cooler upper air, owing to its greater density or weight, settles to the earth. It will be noted that clouds not only prevent the escape of the warm air in immediate contact with the earth, but also blanket, as it were, the upper part of the lower air stratum. Calm or comparatively still air is a condition which favors the formation of frost. On windy nights the air is disturbed and is not permitted to arrange itself in layers according to its density, with the densest and coldest air near the surface of the earth; it is kept mixed up by the wind.

A very interesting fact, which illustrates the manner in which cold air settles by its greater weight to the surface of the earth and allows the warmer air to assume higher levels, causing frost in low grounds and granting to the higher grounds a comparative immunity from frost, is set forth in the following article, by Prof. Cleveland Abbe, in the Monthly Weather Review for December, 1893, entitled Thermal Belts, Frostless Belts, or Verdant Zones:

These are local names given to certain regions on mountain sides within which nocturnal frost rarely or never occurs in the springtime, although freezing tempera-

tures occur in the winter time. Consequently tender vegetation flourishes with remarkable vigor in these regions. The following are the only references to thermal belts in the United States at present known to the editor, but as such regions are specially important to the horticulturist and agriculturist it is hoped that the correspondents of the Weather Bureau will bring these thermal belts to notice wherever they occur in order that their meteorological peculiarities may be better understood.

In the Agricultural Report of the Patent Office for 1861 Mr. Silas McDowell, of Franklin, Macon County, N. C., describes the verdant zone in that county in the valley of the Little Tennessee River. It occupies the region between 300 and 700 feet above the valley of the river, which latter is about 2,000 feet above sea level. On tracing this zone up among the smaller tributaries of the Tennessee River he found that in the higher valleys, where the bottom land is about 3,900 feet above sea level, the verdant zone lies between 4,000 and 4,100 feet. Within this zone frost never injures the vegetation, and the most tender grapes never fail to produce abundant crops.

Prof. J. W. Chickering, jr., in the bulletin of the Philosophical Society of Washington, March, 1883, and in the American Meteorological Journal, volume 1, describes the following thermal belt: "In Polk County, N. C., along the eastern slope of the Tryon Mountain range, in latitude north 35, the thermal belt begins at the base of the mountain, at an elevation of 1,200 feet above the sea, and extends up 2,200 feet, being most perfect at about 1,500 feet. It is about 8 miles long, and is distinguished by magnificent flora, such as would be characteristic of a point 3 degrees south of the actual latitude."

Prof. John Leconte, of Berkeley, Cal., in Science, volume 1, page 278, states that at Flat Rock, near Hendersonville, Henderson County, N. C., on the flank of the mountain spur adjacent to the valleys of the Blue Ridge, he also observed a frostless zone. The valley is about 2,200 feet above sea level, and the thermal belt is 200 to 300 feet above the valley.

In the American Meteorological Journal, volume 1, Mr. S. Alexander describes a thermal belt in which the peach tree flourishes in the southeastern portion of Michigan. He shows that the cold island discovered by Winchell in that region is really the bottom of a topographical depression into which the cold air settles. It is a long valley surrounded by a belt of elevated country from 50 to 600 feet above lakes Michigan and Huron. The valley and the isotherms trend northeast and southwest from Huron County through Sanilac, Lapeer, Oakland, Livingston, and Washtenaw to Hillsdale County. The highlands of this region are all much freer from frost than the lowlands, and all much more favorable for early vegetation. He does not state that any point is high enough to be above the thermal belt, but that, in general, two equal parallel thermal belts inclose the cold island between them.

The thermal belts of California are described in various publications of the Weather Bureau, particularly Bulletin L, "Climatology of California."

In the various numbers of the Monthly Weather Review and in the various summaries published by section directors and district editors special attention has been given to zones of little injury from frost. In nearly every section there are some specially favored localities. One example will serve to illustrate. In the Monthly Weather Review for May, 1910, Professor Howard, secretary of the Missouri state board of horticulture, calls attention to the fact that certain sections in that State, for no apparent reasons based upon

topography or isotherms, nearly always escape injury. He instances particularly the famous peach region about the town of Koshkonong, in Oregon County. Mr. George Reeder, section director, after a careful examination of climatic data, comes to the conclusion that the apparent immunity of orchards in the neighborhood under discussion from damaging temperatures in the spring is probably due to local air drainage, the cold air being drained away from the orchards. He gives tables showing that the cold air spreads out and is drained away down the southern slope of the Ozark plateau, sometimes more toward the southwestward, then toward the south, and again the flow may affect only the southeast. Mr. Reeder states:

It appears as if the cold air current had divided, one part rolling toward the White River Valley, as indicated by the temperature at Hollister, and the other part going toward the Black River Valley, as indicated by the temperature at Doniphan. Instead of the favorable peach belt, if any, extending east and west as suggested by Professor Howard, it probably extends northerly and southerly, or, in other words, it lies along the Ozark border, a narrow strip of table land having elevations ranging between 1,000 feet, on which are situated Koshkonong, Mo., and Mammoth Springs, Hardy, and La Crosse, Ark. This strip bends not far from La Crosse and thence extends northwestward above the White River Valley. The temperature on this border does not fall so low at times as does that at lower as well as higher altitudes.

The cold of April 25 and 26, 1910, that caused so much damage to the fruit crop of Missouri, was quite uniformly distributed over the section under discussion, as may be seen from the following table:

County.	Station.	Temperature.	Date.
		° F.	
McDonald	Dean	28	Apr. 26
Taney	Hollister	31	Apr. 26
Howell	Olden	27	Apr. 25
Oregon	Koshkonong	29	Apr. 25
Ripley	Doniphan	32	Apr. 25

The cold spell was accompanied by cloudy skies and rain mixed with snow, and the movement of the cold currents of air was quite different from that which takes place under a clear sky, or when there is no rain or snow. It will be observed that during cold periods, with more or less cloudiness and moisture, the temperature on the hills may fall to a lower value than the temperature in the valleys; on the other hand, had clear weather prevailed during the nights of April 25 and 26, the temperature conditions would have been reversed, the valleys probably being much colder. It is also quite clear, had the fruit in the valleys escaped winter injury it would have received no more damage from the freeze of April than was sustained in the neighborhood of Koshkonong.

Professor Abbe states in Monthly Weather Review for December, 1903:

It is generally conceded that these thermal belts depend both upon the drainage of cold air downward into the lower valleys and the freedom of radiation from the surface of the ground to the clear sky overhead. During a still night, when

frosts occur, the surface of the hillside cools by radiation, and hence cools the air in contact with it; the latter flows downward as long as its cooling by radiation and conduction exceeds its warming by compression. Inasmuch as its cooling depends on contact with a still colder soil or plant, it soon accumulates in the lowlands as a layer of cold air, which grows thicker during the night by the steady addition of the thin layer of descending air in contact with the ground on the hillsides. The warmer air, which has not yet had an opportunity to cool by contact with the ground, floats on top of the cold mass; it spreads out toward the hills, and is continuously furnishing its heat to the adjacent hillsides as fast as it comes in contact with them before it also cools and descends. The formation of the thermal belt seems to depend largely upon this gentle circulation during the night-time. The lower limit of the belt is defined by the depth of the accumulation of cold air in the confined valley and rises higher in proportion as the night is clearer and longer, and also in proportion as the valley is more or less perfectly inclosed. The upper limit of the thermal belt may depend upon the strength of the wind and the general temperature of the air. But if there be no wind, then it depends equally on the freedom of radiation to the clear sky and on the above-described circulation of air.

The facts quoted above show that frost may be expected on low grounds at times when higher grounds escape the visitations, and the lesson they teach is that early and tender crops should, so far as may be practicable, be confined to crests, hillsides, and mountain sides, and later and hardier crops to the lowlands and valleys.

Local climate, as it is influenced and regulated by the proximity of bodies of water, must be given great weight in calculations regarding frost. Frost is less likely to occur in localities swept by moisture-laden air which has crossed a considerable body of water. This is more especially the case in the fall of the year, when the temperature of bodies of water is reduced very slowly; and in the South, where the water temperature continues relatively high throughout the year, the influence of the water is especially marked. During the colder months air is not only warmed in crossing considerable bodies of water, but it also absorbs moisture, which, although invisible in the form of water vapor, has the effect of retarding the radiation of heat from the earth. In fact, so pronounced is the influence of water and water vapor that localities which with reference to exposure to west and northwest winds are, as it were, in the lee of large bodies of water are comparatively free from frost visitations. On the keys of the east Florida coast, which are protected on the west by lagoons, rivers, lakes or bays, frost is said to be unknown, and throughout the central and northern sections of the United States, localities which are protected on the west and northwest by bodies of water of considerable size are not only favored, by reason of their position, with conditions which do not promote the formation of frost, but also receive the benefits derived from heavier falls of snow.

Local soil conditions constitute, to a certain extent, a factor in the formation of frost. As a rapid loss of heat is promoted by an active

evaporation of moisture, it is evident that, with other conditions equal, frost is more likely to occur on damp than on dry ground, provided, however, that the ground is not too moist, for in the latter event the amount of moisture evaporated and added to the air would have a tendency to retard the radiation of heat from the earth. And herein lies the distinction, so far as frost formation is concerned, between moist air and moist soil. An excess of moisture in the air, in preventing, to a degree, the radiation of heat, is unfavorable to the formation of frost. As frost is the moisture of the air in immediate contact with the earth condensed at freezing temperature, it follows that the earth, from which the moisture of the air is drawn, must contain more or less water, and it is evident that damaging frost will occur with a limited rather than with a large quantity of moisture arising from the earth and a moderately moist and comparatively still air. Both of these conditions usually obtain following, but not too closely following, the rains of spring and autumn. The character of the soil as regards its capacity for retaining moisture must also be taken into account, more especially during periods of comparatively dry weather. Moist soil, as well as the vegetation which springs from it, is chilled to a lower temperature by the evaporation of moisture, and is therefore more subject to visitations of frost than soil which is dry, or into which water penetrates deeply, leaving the surface dry, or soil which sheds the rain and does not absorb and hold water. It is important to note, however, that very moist soil, or soil which contains a large amount of surface water, is, owing to the quantity of moisture yielded to the air, not so subject to the heavy and damaging frosts which visit plants on moderately damp ground or ground that has a small capacity for moisture.

INFLUENCE OF SOIL AND VEGETATION ON MINIMUM TEMPERATURE.

Prof. W. L. Moore, while engaged in studying frost formation in the cranberry bogs of Wisconsin, in 1891 to 1894, came to the conclusion that for a given area the occurrence of frost depended more upon the character of the surface, its covering, and the degree of heat and moisture to which it had been subjected for several days preceding than to the general temperature and pressure of the air, one field receiving an injurious frost, another a light frost, and still another none at all, while apparently the same weather conditions prevailed at a moderate distance above the ground. Experiments carried on since 1906 by Professor Cox in connection with the protection of cranberries show that in the bogs there may be surface differences of from 5° to 10° in temperature, while at an elevation

of 3 feet these differences disappear. Professor Cox is of opinion that variation in temperature is due largely to differences in temperature of the soil or covering. It is as if heaters of varying power were scattered over the bog, giving off heat to the air immediately above. Investigations carried on at Berlin, Cranmoor, and Mather show that in sections which were carefully weeded and cleaned and were free from moss the temperatures were higher. Progressive cranberry growers find that by cultivating, draining, and sanding, low night temperatures are prevented.

DATES OF EARLY AUTUMN AND LATE SPRING FROSTS.

The following table shows for stations of the Weather Bureau throughout the United States dates of first killing frost in autumn and of last killing frost in spring; also the average dates of first killing frost in autumn and of last killing frost in spring:

Dates of first killing frost in autumn and of last killing frost in spring at Weather Bureau stations throughout the United States.

Station.	Date of earliest recorded killing frost in autumn.	Date of latest recorded killing frost in spring.	Average date of first killing frost in autumn.	Average date of last killing frost in spring.
Eastport, Me.	Sept. 8	June 19	Oct. 12	Apr. 28
Northfield, Vt.	Aug. 27	June 7	Sept. 16	May 13
Portland, Me.	Sept. 11	May 31	Oct. 18	Apr. 14
Boston, Mass.	Sept. 30	May 11	Oct. 22	Apr. 26
Nantucket, Mass.	Oct. 1	Apr. 24	Nov. 5	Apr. 10
Block Island, R. I.	Oct. 30	May 11	Nov. 16	Apr. 12
Narragansett, R. I.	Sept. 21	May 30	Oct. 16	Apr. 23
New Haven, Conn.	Sept. 15	...do.....	Oct. 17	Apr. 20
Binghamton, N. Y.	Sept. 16	May 21	Oct. 6	Apr. 27
Albany, N. Y.	Sept. 23	May 30	Oct. 17	Apr. 24
New York, N. Y.	Oct. 5	Apr. 30	Nov. 6	Apr. 10
Harrisburg, Pa.	Oct. 3	Apr. 26	Oct. 24	Do.
Philadelphia, Pa.	...do.....	Apr. 29	Oct. 30	Apr. 8
Atlantic City, N. J.	Oct. 1	Apr. 25	Nov. 4	Apr. 11
Baltimore, Md.	Oct. 6	May 3	...do.....	Apr. 4
Washington, D. C.	Oct. 2	May 11	Oct. 21	Apr. 7
Lynchburg, Va.	Oct. 4	May 7	Nov. 1	Apr. 14
Norfolk, Va.	Oct. 15	Apr. 26	Nov. 12	Mar. 27
Wytheville, Va.	Sept. 14	May 26	Oct. 7	Apr. 23
Charlotte, N. C.	Oct. 8	Apr. 26	Nov. 4	Apr. 1
Raleigh, N. C.	...do.....	May 6	...do.....	Apr. 5
Hatteras, N. C.	Nov. 7	Apr. 19	Dec. 11	Feb. 28
Wilmington, N. C.	Oct. 16	May 1	Nov. 15	Mar. 27
Charleston, S. C.	Nov. 9	Apr. 2	Nov. 30	Mar. 1
Columbia, S. C.	Oct. 19	Apr. 17	Nov. 8	Mar. 23
Augusta, Ga.	Oct. 8	...do.....	Nov. 9	Mar. 20
Savannah, Ga.	Nov. 1	Apr. 5	Nov. 26	Feb. 27
Jacksonville, Fla.	Nov. 12	Apr. 6	Dec. 6	Feb. 14
Jupiter, Fla.	Nov. 18	Apr. 7	Dec. 29	Do.
Atlanta, Ga.	Oct. 11	Apr. 15	Nov. 7	Mar. 24
Macon, Ga.	Nov. 5	Apr. 18	Nov. 14	Mar. 15
Tampa, Fla.	Nov. 28	Mar. 19	Jan. 9	Feb. 8
Pensacola, Fla.	Nov. 12	Apr. 6	Dec. 5	Feb. 23
Mobile, Ala.	Oct. 31	Mar. 28	Nov. 30	Feb. 24
Montgomery, Ala.	Oct. 21	Apr. 5	Nov. 8	Mar. 11
Meridian, Miss.	Oct. 8	Apr. 10	Nov. 2	Mar. 20
Vicksburg, Miss.	Oct. 19	Apr. 6	Nov. 12	Mar. 6
New Orleans, La.	Nov. 19	Mar. 27	Dec. 15	Jan. 24
Shreveport, La.	Oct. 20	Apr. 2	Nov. 11	Mar. 4
Fort Smith, Ark.	Oct. 15	Apr. 6	Nov. 4	Do.
Little Rock, Ark.	Oct. 22	Apr. 14	Nov. 10	Mar. 20

Dates of first killing frost in autumn and of last killing frost in spring at Weather Bureau stations throughout the United States—Continued.

Station.	Date of earliest recorded killing frost in autumn.	Date of latest recorded killing frost in spring.	Average date of first killing frost in autumn.	Average date of last killing frost in spring.
Palestine, Tex.	Oct. 20	May 30	Nov. 13	Mar. 13
Galveston, Tex.	Dec. 4	Mar. 1	Dec. 25	Feb. 5
San Antonio, Tex.	Nov. 9	Mar. 20	Nov. 30	Feb. 25
Corpus Christi, Tex.	Nov. 30	Mar. 19	Dec. 25	Feb. 27
Memphis, Tenn.	Oct. 2	Apr. 16	Oct. 28	Mar. 24
Nashville, Tenn.	Oct. 8	May 14	Oct. 24	Apr. 2
Chattanooga, Tenn.	Sept. 30	...do....	Oct. 26	Do.
Knoxville, Tenn.	Oct. 1	Apr. 24	Oct. 27	Apr. 3
Louisville, Ky.	Sept. 24	May 14	Oct. 29	Apr. 6
Lexington, Ky.	Sept. 30	May 20	Oct. 23	Apr. 18
Evansville, Ind.	...do....	Apr. 21	Oct. 30	Apr. 10
Indianapolis, Ind.	Sept. 21	May 21	Oct. 19	Apr. 16
Cincinnati, Ohio.	Sept. 30	Apr. 24	Oct. 25	Apr. 14
Columbus, Ohio.	Sept. 21	May 17	Oct. 16	Apr. 16
Elkins, W. Va.	Sept. 28	May 24	Oct. 10	May 18
Parkersburg, W. Va.	Sept. 24	May 22	Oct. 17	Apr. 11
Pittsburg, Pa.	Sept. 25	May 29	Oct. 19	Apr. 26
Oswego, N. Y.	...do....	...do....	...do....	Apr. 27
Rochester, N. Y.	Sept. 26	May 30	...do....	May 1
Buffalo, N. Y.	Sept. 23	May 29	Oct. 16	Apr. 25
Erie, Pa.	Oct. 12	May 17	Oct. 29	Apr. 22
Cleveland, Ohio.	Oct. 2	May 22	Oct. 31	Apr. 16
Sandusky, Ohio.	Oct. 8	May 17	Oct. 26	Apr. 14
Toledo, Ohio.	Sept. 9	May 29	Oct. 15	Apr. 24
Detroit, Mich.	Sept. 21	May 31	Oct. 11	Apr. 30
Port Huron, Mich.	Sept. 22	June 6	Oct. 9	May 8
Alpena, Mich.	Sept. 6	June 9	Sept. 26	May 14
Sault Ste. Marie, Mich.	Sept. 5	May 29	Sept. 24	May 16
Marquette, Mich.	Aug. 22	June 11	Oct. 2	May 15
Escanaba, Mich.	Sept. 9	June 16	Oct. 1	May 14
Green Bay, Wis.	Sept. 16	May 30	Oct. 4	May 5
Grand Haven, Mich.	Sept. 23	May 28	Oct. 10	Apr. 28
Milwaukee, Wis.	Sept. 25	May 29	...do....	Apr. 29
Chicago, Ill.	Sept. 18	...do....	Oct. 15	Apr. 18
Duluth, Minn.	Sept. 15	June 8	Oct. 5	May 3
Minnespolis, Minn.	Sept. 13	May 20	Oct. 8	Apr. 27
St. Paul, Minn.	Sept. 20	May 25	Oct. 5	May 6
Madison, Wis.	Sept. 29	May 13	Oct. 17	Apr. 21
La Crosse, Wis.	Sept. 21	June 1	Oct. 8	May 2
Dubuque, Iowa.	Sept. 27	May 21	Oct. 13	Apr. 20
Davenport, Iowa.	Sept. 18	May 22	Oct. 14	Apr. 22
Des Moines, Iowa.	Sept. 12	...do....	Oct. 8	Apr. 28
Keokuk, Iowa.	Sept. 18	May 4	Oct. 15	Apr. 13
Hannibal, Mo.	Sept. 30	May 14	...do....	Apr. 14
Springfield, Ill.	Sept. 25	May 22	Oct. 16	Apr. 20
St. Louis, Mo.	Sept. 30	...do....	Oct. 27	Apr. 2
Cairo, Ill.	...do....	Apr. 19	Oct. 28	Mar. 29
Springfield, Mo.	...do....	May 19	Oct. 13	Apr. 16
Columbia, Mo.	Sept. 28	May 9	Oct. 15	Apr. 14
Kansas City, Mo.	Sept. 30	May 4	Oct. 24	Apr. 10
Topeka, Kans.	Sept. 28	May 19	Oct. 13	Apr. 8
Wichita, Kans.	Sept. 23	May 15	Oct. 19	Apr. 6
Concordia, Kans.	Sept. 27	May 19	Oct. 14	Apr. 24
Omaha, Nebr.	Sept. 18	...do....	Oct. 12	Apr. 15
Lincoln, Nebr.	Sept. 12	May 10	Oct. 9	Apr. 18
Valentine, Nebr.	...do....	June 21	Sept. 18	May 9
Sioux City, Iowa.	Sept. 13	May 21	Sept. 27	May 1
Huron, S. Dak.	Aug. 23	June 8	Sept. 20	May 13
Pierre, S. Dak.	Sept. 12	May 19	Sept. 30	Apr. 30
Moorhead, Minn.	Aug. 25	June 8	Sept. 22	May 14
Bismarck, N. Dak.	Aug. 23	June 7	Sept. 21	May 15
Williston, N. Dak.	Aug. 18	June 10	Sept. 14	May 18
Havre, Mont.	Aug. 27	June 6	Sept. 13	May 14
Helena, Mont.	Sept. 5	June 9	Sept. 25	May 10
Rapid City, S. Dak.	Sept. 13	May 21	Sept. 20	May 1
Lander, Wyo.	Aug. 23	June 18	Sept. 11	May 19
Salt Lake City, Utah.	Sept. 22	...do....	Oct. 18	Apr. 23
Grand Junction, Colo.	Sept. 27	May 15	Oct. 21	Apr. 14
Cheyenne, Wyo.	Aug. 29	June 11	Sept. 16	May 22
North Platte, Nebr.	Sept. 10	May 23	Sept. 28	May 1
Denver, Colo.	Sept. 12	June 6	Oct. 5	May 6
Amarillo, Tex.	Oct. 16	May 23	Nov. 1	Apr. 16
Pueblo, Colo.	Sept. 12	...do....	Oct. 5	Apr. 26
Dodge, Kans.	Sept. 23	May 27	Oct. 15	Apr. 17

Dates of first killing frost in autumn and of last killing frost in spring at Weather Bureau stations throughout the United States—Continued.

Station.	Date of earliest recorded killing frost in autumn.	Date of latest recorded killing frost in spring.	Average date of first killing frost in autumn.	Average date of last killing frost in spring.
Oklahoma, Okla	Oct. 7	Apr. 30	Oct. 31	Apr. 2
Fort Worth, Tex	Oct. 22	May 1	Nov. 19	Mar. 18
Abilene, Tex	Oct. 24	Apr. 16	Nov. 15	Mar. 15
El Paso, Tex	Oct. 23	Apr. 22	Nov. 10	Mar. 20
Santa Fe, N. Mex	Sept. 27	May 18	Oct. 19	Apr. 15
Yuma, Ariz	None.	None.	None.	None.
Phoenix, Ariz	Nov. 9	Mar. 31	Dec. 7	Feb. 18
Spokane, Wash	Sept. 7	June 8	Oct. 17	Mar. 23
Walla Walla, Wash	Sept. 28	May 12	Nov. 7	Apr. 4
Seattle, Wash	Oct. 18	May 27	Nov. 22	Mar. 21
Portland, Oreg	Oct. 13	May 9	Nov. 16	Mar. 17
Tatoosh Island, Wash	Nov. 1	Apr. 19	Dec. 15	Mar. 14
Roseburg, Oreg	Sept. 25	May 16	Oct. 30	Apr. 15
Baker City, Oreg	Sept. 4	June 24	Sept. 27	May 25
Winnemucca, Nev	Aug. 22	June 20	Sept. 24	May 15
Eureka, Cal	Nov. 7	May 1	Dec. 8	Apr. 2
Red Bluff, Cal	Nov. 15	Apr. 25	Dec. 6	Feb. 26
Sacramento, Cal	Oct. 28	Apr. 26	Nov. 15	Feb. 16
San Francisco, Cal	Nov. 18	Apr. 20	Dec. 26	Jan. 8
Fresno, Cal	...do....	Apr. 14	Dec. 15	Mar. 4
San Luis Obispo, Cal	Nov. 19	Apr. 8	Dec. 27	Feb. 20
Los Angeles, Cal	None.	...do.....	None.	Apr. 8
San Diego, Cal	None.	None.	None.	None.

PROTECTION FROM FROST.

A discussion of methods of protection from frost calls for a consideration in great detail, not only of the several approved processes of protection and the extent to which they can be applied in connection with the various crops, but also of the value of these processes under the varying conditions of local topography, local climate, and local soil conditions.

It is evident, in the light of the foregoing statements, that fruit and vegetable growers in rolling and hilly country, or when located in the lee, as regards west to northwest winds, of bodies of water, can, to some extent, so place in their earlier and tenderer fruits and crops that they will, in the first instance, avoid the lowlands and valleys, and in the second take advantage of the more moderate and moist air that crosses water surfaces. For further protection the grower must depend upon artificial appliances. These appliances are designed to produce the following effects or results: To prevent a rapid radiation of heat from the earth; to charge the air with moisture; to warm the air; to create artificial drafts, whereby the air is mixed and the cold air is not allowed to settle to the surface of the earth; or to actually cover or roof in plants.

DEVICES FOR PREVENTING RAPID RADIATION OF HEAT.

Devices designed to prevent a rapid radiation of heat from the earth include screens which can be drawn over plants, vineyards, and groves; loose substances with which low plants may be covered; and

smudge fires built to the windward of areas for which protection is desired. These devices, which of a necessity are operative for very limited areas, have been described by Prof. W. H. Hammon and by Prof. A. G. McAdie.

Glass screens.—In greenhouses and hotbeds advantage is taken of the peculiar property of glass which allows the heat rays of the sun to pass through it, and is almost impervious to the dark *heat* rays from the earth and plants. This is one of the most perfect screens possible, since it not only prevents the loss of heat by radiation, but receives and retains the heat from the sun. The expense precludes its adoption except for the protection of valuable plants and flowers.

Screens of other solid materials have been quite extensively used in protecting vineyards and citrus groves where intense cultivation is practiced and where the location of the groves near an excellent market admits of profit even with expensive methods of cultivation.

Cloth screens.—In Italy and portions of France screens made of muslin strung on wires stretched on poles above the tops of trees or vines have been used extensively. These screens are drawn on nights when frosts are probable, and are pushed back during the day. When the season has advanced so far as to preclude further danger, they are taken down and stored. Of course such a plan could be operated only on a very limited scale, and would then be expensive. This plan has been recently successfully tried in the orange groves of southern California.

Lath screens.—During the past few years screens made of laths fastened to ordinary telephone wire (the spaces between them being about the width of the laths) have been extensively used in Florida. These are spread over a frame erected above the trees or plants. The screens serve not only as a fair protection from frost, but also as a shade from the hot sun. When no longer needed, they can be rolled up and stored away for preservation. At first thought it would seem improbable that a screen covering only half the space (the spaces being as wide as the laths) would afford much protection; but when it is considered that laths have considerable thickness it is plain that while only one-half the vertical rays are screened those inclined between the vertical and horizontal are partially intercepted by the edges as well as the faces of the laths. As a matter of fact, about three-fourths of the sky is screened by this means. By placing the laths in north and south directions the direct rays of the morning sun are completely cut off from the orchard, which admits of the temperature rising slowly. This reduces the liability of injury to plants.

Paper cover.—A new form of cover, known as an antifrost cover, has been devised in the San Francisco Weather Bureau office. It is, however, best suited for the protection of small fruit, garden flowers, and vegetables. It is an old and widely known practice to cover garden flowers with newspapers when frost is expected. In the antifrost cover there is used instead of a single cover a double layer of prepared paper with an intervening air space. This prevents almost perfectly the escape of the long heat waves from the ground. Such covering can be spread above the plant an hour or two before sunset, i. e., before the ground has lost much of its heat. If at the same time shallow pans of warm water are placed under the cover, an effective screen and serviceable supply of water are provided. It is also to be noted that we use this cover at a time when the roots are absorbing vigorously and transpiration by the leaves is at a maximum. We therefore prevent any lowering of temperature at the leaf surface and store up in the plant a quantity of heat energy for expenditure slowly through the night hours. The actual temperature of the air is of less importance than the temperature of the leaf or fruit surface. A deposit of moisture is advantageous.

A special use of the antifrost cover in connection with tree fruits is to unroll the covers in the orchard before sunset and to roll them up about 4 o'clock in the morning.

Other methods.—Strawberries and other low plants are frequently protected by covering them with straw or other loose substances.

Frequently young potato plants are saved by plowing a furrow alongside and allowing the dirt to bury them.

Cranberry growers in the marshes of Massachusetts, New Jersey, and Wisconsin flood the marshes when frost is expected. In this case the protection is probably due, for the most part, to the high specific heat of the water, as only portions of this land are submerged. Cultivating, draining, and sanding are now resorted to with such good results that there is practically no need of flooding except in spring and autumn.

DEVICES FOR ADDING HEAT TO THE AIR.

The first successful attempts to prevent injury by frost by the use of heating devices on a large scale occurred in California about 1896. Among a number of protective methods based upon heating, described by Prof. A. G. McAdie, may be mentioned the device of Edward Copely, of Riverside, Cal., in which a wire basket filled with coal and sufficient kindling to insure proper combustion was used. The method met with much favor in the orange groves of California. With from 20 to 40 of these baskets to the acre the temperature under ordinary winter weather conditions can be raised 3 degrees or more.

The first use of oil that we have a record of was by Everett, at Arlington, Cal., and the first use of hot water was by Meacham at Riverside.

A severe frost about the end of December, 1895, caused great damage to the orange crop in the Riverside section, and the following year may be said to have marked the beginning of the frost-fighting campaign. The Riverside Horticultural Club took an active interest in the problem and many experiments were conducted by such men as those above mentioned and Koethen, Reed, Holmes, Hall, Hammon, McAdie, and others. The work has gone on, and there are now on the market many forms of heating devices. A convenient and serviceable frost alarm thermometer and oil pot were devised by the Weather Bureau official at Fresno, Cal., Mr. J. P. Bolton, for use in the vineyards during the spring frosts. There have recently been put upon the market in the apple sections of Colorado and other Rocky Mountain States orchard heaters burning oil, coal, or other fuel.

The various forms of heaters are known by the name of their inventors or are otherwise designated. It is not desirable in a bulletin of this character to advocate the use of any one special form. Indeed the heater that is most satisfactory in one locality and for a given crop may prove less satisfactory under changed conditions.

As showing the efficiency of these heaters, the following report upon a certain coal-burning heater was made by Prof. O. B. Whipple, field horticulturist, Colorado Experiment Station:

I have the following report to make on the coal-burning heater demonstration, held at your place on the evening of November 12, 1908:

The conditions under which the test were held were such as one would experience in fighting frost in the spring. One hundred heaters were well spaced at 29.5 feet,

covering 2 acres of ground. In placing the thermometers, I was careful to see that no pots were closer than 20.5 feet. Waste saturated with one-half crude oil and one-half kerosene with tree prunings were used to start the fires. The fires were started at 7.45, and responded promptly, as the table will show. One man lighted the 100 pots in eight minutes. All pots started without trouble. The pots were refilled, beginning at 10.15, one man completing the job in forty minutes. A few of the pots on a part of the block were again refilled at 12.30, and all burned until the end of the test. In all 2 tons of coal were consumed in the seven hours, making a total of 40 pounds per pot. The coal used was one-half lump and one-half nut and came from the Farmers' mine and Stokes' mine. The temperature readings shown in the table were taken at an elevation of 7 feet from the ground. A thermometer at 12 feet from the ground inside the heated area showed one degree higher temperature.

The following table shows the temperature readings for the full seven hours:

Hour.	Outside temperature.	Inside temperature.	Gain.	Hour.	Outside temperature.	Inside temperature.	Gain.
p. m.	°*F.*	°*F.*	°*F.*	*p. m.*	°*F.*	°*F.*	°*F.*
7.45	30.2	30.2	.....	12.00	29.3	36.9	7.6
8.15	31.1	35.6	4.5	*a. m.*			
8.30	28.4	32.9	4.5	12.30	25.7	32.6	69.
9.00	28.4	34.2	5.8	1.00	26.1	32.4	6.3
9.30	28.4	33.8	5.4	1.30	26.6	33.8	7.2
10.00	28.4	34.2	5.8	2.00	28.4	35.6	7.2
10.30	27.5	32.9	5.4	2.30	28.4	34.5	6.1
11.00	27.5	33.8	6.3	3.00	28.4	33.4	5.0
11.30	28.4	36.0	7.6				

The following table is from the report of the committee in the "under actual frost conditions" test held at W. C. Knight's orchard, Grand Junction, Colo., October 28, 1908. The committee consisted of O. B. Whipple, field horticulturist Colorado Experiment Station; W. C. Knight and John M. Harvey, fruitgrowers.

Outside temperature.	Hour.	Inside temperature, with 50 coal heaters with concentrated draft.	Gain.	Inside temperature, with 100 oil heaters, best make.	Gain.	Inside temperature, with 80 coal heaters, open draft on all sides.	Gain.
°*F.*	*p. m.*	°*F.*	°*F.*	°*F.*	°*F.*	°*F.*	°*F.*
35	8.00	35	.....	35	.....	35	.....
35	8.15	38	3	38	3	36	.....
33	8.30	39	6	39	6	38	5
32	8.45	40	8	40	8	37	5
34	9.00	40	6	39	5	38	4
34	9.30	40	6	39	5	37	3
34	10.00	40	6	38	4	38	4
32	10.30	36	4	37	5	36½	4½
31	11.00	35	4	36	5	35	4
30	11.30	36½	6½	36	6	34	4
28	12.00	35¼	7¼	36	6	34	5
	a. m.						
27	12.30	34½	7½	35	8	32	5
27	1.00	34½	7½	35	8	32	5
Average gain.....			6	.....	5¾	.....	4.4

The following account of the use of smudge pots, as they are called, is given by Mr. Richard H. Sullivan, local forecaster, in the Monthly Weather Review for 1910, vol. 38, pp. 412–413:

The inclosed series of observations were taken for the benefit of members of the Sedgwick County Horticultural Society and others in connection with the frost warnings of March 30 and April 5 and tests of smudge pots in the 1-acre orchard of Mr. Albert Kunkel, in Wichita, on March 31 and April 6, and in a selected plat of the orchard of the Thomas Orchard Company, 3 miles west of Wichita, on April 6. The readings for the Kunkel orchard on March 31 and the Thomas orchard on April 6 were made by the writer, 2 minimum and 5 exposed thermometers, 1 anemometer, and 1 commercial thermometer being used.

On Mr. Kunkel's place 79 pots to the acre were used on March 31 and 70 pots to the acre on April 6; his fruit is still unharmed. In the Thomas orchard, 50 pots to the acre, or 500 pots in all, were used among Jonathan and Grimes Golden apple trees. Unfortunately, the temperature in the latter orchard could not be kept above freezing after 4 a. m. on account of lack of fuel oil. The small number of pots to the acre made it necessary to run them at nearly full capacity, and in order to carry the heat through to sunrise the lighting was delayed until 2 a. m. However, the manager does not consider his loss as great as in some orchards in the vicinity.

The figures show that with from 70 to 80 pots to the acre a fruit crop can be saved when the temperature falls to 25°, or even to 22°, if the work is done thoroughly and systematically. These are the first known practical tests of this character that have been made in this vicinity, and the whole proposition rests upon the question of how much expense for such insurance the investment can stand and still render a profit.

Where artificial methods of prevention of damage were not used during the freezes of March 31 and April 6, especially in the lowlands, there was general loss of apricots, peaches, plums, pears, and a very large proportion of apple buds, making the third series of disastrous spring frosts in 4 years. Last year the whole fruit-tree crop was killed by a single freeze during the night of April 30–May 1, warning of which was given the day before.

There is now a general disposition among local horticulturists to adopt the methods of commercial fruit growers elsewhere to prevent damage by frosts and freezes, and the belief is steadily growing that the warnings of this service must be heeded if loss is to be avoided.

Comparative temperature, wind velocity, and weather readings made in connection with test of central-draft smudge pots for prevention of damage by frost or freezing in the orchard of Albert Kunkel.

Half-hourly observations were taken from midnight of March 30–31 to 7.30 a. m., March 31. The arrangement and height of the various instruments are indicated by the letters in the diagram (fig. 3) and the explanation herewith: *A*, Weather Bureau anemometer, 5 feet above ground; *a*, W. B. minimum thermometer No. 6733, 3 feet 8 inches above ground; *b*, W. B. standard thermometer No. 4403, 4 feet 9 inches above ground; *c*, W. B. standard thermometer No. 4788, 4 feet 9 inches above ground; *d*, W. B. standard thermometer No. 4789, 4 feet 9 inches above ground; *e*, W. B. minimum thermometer No. 9436, 3 feet 8 inches above ground; *f*, W. B. standard thermometer No. 3135, 4 feet 9 inches above ground; *g*, W. B. standard thermometer No. 5671, 4 feet 9 inches above ground; *h*, commercial thermometer, 1 foot above ground; *i*, residence; *j*, greenhouse; *k*, barn.

The smudge pots, 79 in number, were lighted between 5.30 and 5.45 a. m., March 31, and extinguished at 7.30 a. m. The temperatures at 6 a. m. in the accompanying table show the effectiveness of such a method during frosty periods. Fuel oil, costing 4 cents per gallon, was used.



FIG. 3.—Diagram of Kunkel orchard, Wichita, Kans. Shaded areas show limits of smudge pots.

Winds—Northwesterly to 2.30 a. m.; northerly to 6.30 a. m.; northeasterly to 7.30 a. m.

TABLE 1.—*Temperature, weather, and wind velocity, March 31, 1910.*

[Letters in heading indicate position of thermometers.]

Time.	*a*	*b*	*c*	*d*	*e*	*f*	*g*	*h*	Weather.	*A*
	°F.	°F.	°F.	°F.	°F.	°F.	°F.	°F.		Miles
12.00 midnight	39.0	40.5	41.0	41.0	39.0	41.0	40.5	39.0	Clear.	0.2
12.30 a. m	38.5	40.0	39.5	39.5	38.5	39.5	39.5	38.0	Clear.	0.2
1.00 a. m	36.5	38.5	38.0	38.0	37.0	38.5	38.5	37.0	Clear.	0.2
1.30 a. m	35.5	37.5	37.0	37.0	36.0	38.0	37.5	36.0	Clear.	0.2
2.00 a. m	35.0	37.0	37.0	36.0	35.5	36.0	36.0	34.5	Clear.	0.3
2.30 a. m	34.0	36.0	35.5	35.5	34.0	35.5	35.5	34.0	Clear.	0.3
3.00 a. m	35.5	37.0	37.5	37.0	36.0	37.5	37.0	35.0	Clear.	0.4
3.30 a. m	36.0	37.0	39.0	39.0	37.0	38.5	38.0	36.0	Clear.	0.5
4.00 a. m	36.5	37.0	36.5	36.0	35.5	37.0	37.0	36.0	Clear.	0.3
4.30 a. m	35.5	36.5	36.0	36.0	35.0	36.5	36.0	35.0	Clear.	0.3
5.00 a. m	33.0	34.0	33.5	33.0	33.0	34.0	33.0	31.0	Clear.	0.2
5.30 a. m	a31.0	32.5	32.5	32.0	31.5	33.0	33.0	30.0	Clear.	0.2
6.00 a. m	31.0	44.0	40.5	38.0	34.0	38.0	38.0	36.0	Clear.	0.3
6.30 a. m	33.0	39.0	40.0	38.5	34.5	37.5	38.0	38.0	Clear.	0.4
7.0 a. m	35.5	38.0	38.0	37.5	35.5	38.0	39.0	38.0	Clear.	0.5
7.30 a. m	40.0	40.5	41.0	41.0	40.0	40.5	40.5	40.0	Clear.	0.5

a Lowest temperatures recorded, 30.5° between 5.00 and 5.30 a. m.

79 smudge pots lighted, beginning at 5.30 a. m. and ending at 5.45 a. m.

Second test made by Mr. Albert Kunkel. Number of pots lighted, 70.

TABLE 2.—*Temperature and weather table, April 6, 1910.*

[Letters in heading indicate location of thermometers as shown on diagram of March 31, 1910.]

Time.	a	b	c	Weather.
	°F.	°F.	°F.	
3.00 a. m.	30	32	.....	Clear.
3.30 a. m.	28	31	.....	Clear.
4.00 a. m.	27	34	.....	Clear.
4.30 a. m.	26	35	.....	Clear.
5.00 a. m.	26	36	.....	Clear.
5.30 a. m.	27	36	37	Clear.
6.00 a. m.	28	38	38	Clear.
6.30 a. m.	28	35	38	Clear.
7.00 a. m.	30	38	38	Clear.

70 smudge pots lighted, beginning at 3.30 a. m.

Comparative temperature, wind velocity, and weather readings made in connection with test of central-draft smudge pots for prevention of damage by frost or freezing in the orchard of the Thomas Orchard Company, 3 miles west of Wichita.

The smudged portion of the orchard was a selected 10-acre plat containing Jonathan and Grimes Golden apples; the pots were arranged 50 to the acre, or 500 in all. The instruments were placed at a height of 6 feet, except 1 commercial thermometer, which was located 1 foot above the ground near the center. The thermometers are



FIG. 4.—Diagram of plat of Thomas Orchard Company, Wichita, Kans.

arranged in the table in the order of reading. The pots were charged with 3 quarts of fuel oil, and lighting was delayed until 2 a. m. on account of lack of oil. The 500 pots were lighted between 2 and 2.35 a. m., and insufficient oil to hold the fires at reasonable capacity resulted in inability to keep temperature above freezing after 4 a. m., and by daylight practically all of the oil had been consumed.

Winds—Northwesterly to 4.30 a. m.; westerly after 4.30 a. m.

The letters in the diagram (fig. 4) of this plat indicate corresponding instruments in Table 3; numbers, United States Weather Bureau thermometers; *d*, commercial thermometer; *A*, anemometer.

TABLE 3.—*Temperature, weather, and wind velocity.*

[Letters in heading indicate position of thermometers.]

Time.	*a*	*b*	*c*	*d*	*e*	*f*	*g*	*h*	Weather.	*A*
	°F.	°F.	°F.	°F.	°F.	°F.	°F.	°F.		*Miles.*
12.00 midnight	31.5	31.5	32.0	31.5	32.0	31.0	31.0	31.0	Clear.	0.2
1.00 a. m.	31.0	31.0	31.0	32.0	31.5	31.0	31.0	30.5	Clear.	0.2
2.00 a. m.	30.0	[a]30.0	30.0	30.0	30.0	29.5	31.5	29.5	Clear.	0.1
3.00 a. m.	33.5	33.5	33.5	35.0	34.0	33.5	34.5	33.0	Clear.	0.1
4.00 a. m.	32.5	32.0	32.5	32.0	32.5	32.5	33.5	32.0	Clear.	0.1
5.00 a. m.	31.0	31.5	31.0	31.5	30.5	29.0	30.0	30.0	Clear.	0.2
6.00 a. m.	31.0	30.0	30.0	30.0	30.0	29.0	29.0	28.5	Clear.	0.1

[a] Lowest temperature recorded, 29.0° between 1.00 and 2.00 a. m.

500 smudge pots lighted, beginning at 2.00 a. m. and ending at 6.35 a. m.

Additional report, April 30, 1910.

Further trials in the Thomas orchard on April 19, 24–25, fully confirm the efficacy of the firing methods above explained in detail. The wind-velocity records show, as was to be expected, that the most damaging temperatures occur after the wind has died down. Artificial preventive methods can be used effectively despite the oft-repeated arguments that it is too windy in Kansas.

In a paper read before the California Fruit Growers' convention at Watsonville, December, 1909, the district forecaster discusses in some detail the various forms of heating devices, such as open fires, smudge fires, coal baskets, oil pots, orchard heaters, etc., as used in the protection of citrus fruits in that State.

Many small fires are found to be much more effective than one or two large fires. The great mass of experiments made in California orchards show that direct heating of the air by open fires has not been sufficient to prevent injury at times of very low temperatures. A large amount of the heat thus produced is wasted; the efficiency of the method is low. This is illustrated by the following experience of a gentleman who is a close observer, an earnest student of the problem of frost-protection, and one in whom I have the greatest confidence. During the night of December 20–21, 1908, on a certain California ranch, the temperature for fourteen hours ranged between 19° and 24° F. For thirty-six hours the temperature did not rise above 28°. During this night the gentleman referred to burned 15 cords of wood and about 40 tons of wet hay in his efforts to protect his orchard. The relative humidity was low, there was little movement of the air, and he reports that "the smoke rose as straight as a pine tree." At the intersection of two roads in the orchard a large fire was maintained, and 30 feet distant the temperature in an olive tree was observed. Another fire was burning within 20 feet of this tree on a second side, and on yet a third side was a third fire maintained within 25 feet of the same tree. The temperature at the tree, however, remained at about 20° F. from 3 to 8 a. m. of December 21. This was the coldest weather in this locality since 1888. It is evident that in this case a large amount of heat escaped without producing the desired warming effect, i. e., was lost, wasted. It is, of course, well known that the rate of conduction of heat through air is low.

DEVICES FOR ADDING MOISTURE TO THE AIR.

Smudge fires made of damp material have been more successfully used for the purpose of affording protection from frost than those made of tar, crude petroleum, and other dry materials, for the reason

that in addition to creating a dense smoke they add to the atmosphere a considerable amount of evaporated water, which, though invisible, serves to retard the radiation of heat. When any considerable amount of water is thus converted into vapor and is distributed in the air which covers the area to be protected, a portion of it is likely to be condensed by the surrounding cooler air and appear in the form of mist, which acts as an agent to prevent the escape of heat from the earth; and the act of condensation sets free in the air some of the heat that has been expended in the process of evaporating the water contained in the smudge-fire materials. Thus the process is well ordered throughout. In the first place, an undue quantity of heat, whereby undesirable drafts would be created, is not communicated to the air by the fires, and the final result is that in reliberating, by condensation, the water in the air the particles of moisture remain suspended in the air, and some of the heat of the fires is returned to the air in a form which, to a certain degree, warms it without causing currents which would be calculated to disturb the smoke cloud, or pall, and the moist air which overlies the ground and upon whose presence the safety of the plants from frost depends.

Portable smudge fires.—A number of excellent devices have been tried, in which the fires were built upon some vehicle by which they could be moved about the orchard. The advantages of this plan are several.

First. The fire can be moved to the section where most needed, which is generally along the windward side of the orchard.

Second. The loss of heat by an upward draft is almost entirely prevented, since the fire does not remain in one position long enough to establish such a draft. On this account much larger and, consequently, fewer fires, with equal efficiency, are possible.

Third. There is a much more uniform distribution of heat and smoke throughout the orchard.

Aside from the smudge fires, other devices have been adopted to charge the air directly with moisture, some of which are constructed on a plan too elaborate and expensive for general use.

The first operation involved is the evaporation of water in large quantities, either by means of evaporating pans or by the use of boilers with connecting pipes whereby steam can be generated, conducted, and discharged into the air in different parts of a field or orchard.

Spraying and sprinkling plants in times of threatened frost has also been attended with some degree of success. At the Everest ranch, at Riverside, Cal., sprinklers are placed at the top of 50-foot masts, which fill the air with a very fine spray. In nearly every instance the protection against frost by this method has been sufficient.

Sprinkling gardens before sunrise on frosty mornings has proved of decided value.

Secondly, irrigation, which of necessity can be used only to a very limited extent and in certain favored localities, has been found to afford very complete protection from frost in fields, orchards, and vineyards which are equipped with irrigation ditches, and in the cranberry districts of Wisconsin and Cape Cod perfect protection from damage by frost is secured by flooding the cranberry marshes the day before heavy frost occurs in those sections.

In no case should water be allowed to stand in orchards, vineyards, or gardens during the early growing period while danger from frost is possible, for the reason that water unduly promotes the growth of young plants and thereby renders them more susceptible to damage by frost and cold than are the plants whose growth is retarded.

GENERAL OBSERVATIONS AND PRACTICAL RESULTS.

The general campaign of frost fighting, as thus far laid out, covers three main lines:

(1) Accurate advance information of the occurrence of frost.
(2) Applying preventive measures during critical hours.
(3) Preventing sudden warming of chilled fruit, or a too rapid thawing out.

Under the first of these heads great progress has been made in the past five years. Throughout the various fruit sections of the United States, Weather Bureau officials recognize the value of special study of types of frost maps. The work first undertaken in California on a large scale and for a distinct purpose is now an important feature of the forecaster's work at a large number of places. Frosts are found to occur as a consequence of certain movements of low and high pressure areas and the displacement of the lower air incident to the pressure changes. Frost is a matter of air drainage, both on a large scale and on a small scale. In other words, frosts are successfully forecast because we understand the general movements of the lower air and in particular certain local circulations.

The forecaster anticipates a condition in which the lower air strata are quiet, dust free, and very dry. Under such condition there is rapid radiation and loss of heat from plant and ground and a sharp fall in temperature. While the forecaster gives the general conditions favorable for frost, it must be clearly understood that each individual orchardist must study his own problem of air movement in his own locality. These local circulations should be studied in connection with the daily weather map. It is also advisable to have certain instrumental records, such as are given by the thermograph and hygrograph.

Under the second head we have discussed the physical processes operative in the formation of frost and mentioned the various devices, such as smudging outfits, coal baskets, oil pots, orchard heaters, antifrost covers, etc. Opinions will differ as to the relative efficiency of various devices, and some are undoubtedly more serviceable for certain localities than others.

There are three general principles used in frost-protection devices:

(1) Adding heat.
(2) Saving heat.
(3) Mixing or stirring the air.

Under the third head, methods based upon mixing or stirring the air, no special devices that are available and practicable are known. When nature mixes the air—i. e., on windy nights—frost does not occur. It is now known to meteorologists that layers of air of different temperature may lie close to one another without mixing. On nearly every frosty night there is a difference of 6, 8, or 10 degrees in temperature between the ground and the air 10 feet above, the warmer layers being above. Where air is well mixed and there is good ventilation, we seldom find frost.

WORK IN FLORIDA.

Devices used in Florida for the protection of fruit, vegetables, and berries from frost and light freezes conform in character to those tested in the California experiments, in which fires were found to be the most effective means of protection, and appliances for adding moisture to the air were successful only to a slight degree.

As artificial appliances are totally inadequate to add to the atmosphere any very appreciable amount of moisture, it is evident that methods which may have been found ineffective in the dry climate of California would possess value in localities where their office is confined to adding moisture to an already moist atmosphere. The Gulf and South Atlantic Coast States, and in fact the country generally from the Mississippi Valley to the Atlantic seaboard, possess a moist atmosphere, and the fruits and tender vegetables of these districts can, therefore, be the more readily protected from frost by devices which add moisture to the air. As a means of adding moisture to the air, irrigation should be more effective than in California; and in localities where this method can be used protection will be assured except against hard freezes.

Owing to the comparatively inexpensive character of the materials used in damp smudge fires, they seem the best adapted for common use in groves and gardens. Berries and other low plants can be protected with but little expense by coverings of straw and other light materials. In Florida pine needles are successfully used to pro-

tect strawberries. Devices for actually heating the free and open air are expensive, and their value is dependent solely upon a comparatively still air and small, numerous, and well-distributed fires.

In Florida many experiments have been made with a view to adopting devices which will protect citrus trees and fruits from injury during the periods of severe cold which at times visit that section. The fact that these periods of cold are infrequent does not relieve the grower from the necessity of providing for their occurrence. Effective measures of protection necessarily vary with the conditions that confront the grower. Protection can be made effective by tents, coverings of cloth, sheds, and one-half and one-third lath shelters and their modifications, that retain heat from open wood fires and sheet-iron stoves. The expense of these methods is obviously too great to make their employment, except for choice varieties of fruit, commercially profitable.

In this State protection of citrus trees and fruit in the open field is usually afforded by what is known as "firing." This method is effective, except in the presence of severe freezes, provided rain or strong winds do not accompany the cold. Heavy rain will, of course, extinguish the fires, and strong winds will not permit a necessary accumulation of warm air about the trees. When no rain is falling and the wind is light, fires must be built on the windward side of the trees, so that the heat will be carried through the trees, and the number of the fires must be regulated by the degree of cold. In this form of protection the grower must lay in and place his materials for "firing" before the winter season sets in, and must, during the season, be on guard to utilize the materials (usually wood) that he has accumulated. When local indications point to dangerously low temperatures, or when warnings have been received from the Weather Bureau, he should watch his thermometers, exposed in the groves, and be prepared to "fire" with a rush when the mercury sinks to the danger point, 28° F., with a downward tendency. There is no time on the eve of a freeze to collect firing materials, and the grower must not desert his grove for bed or board so long as danger by cold continues. He will have his hands full keeping his fires going, even if he has sufficient material for that purpose. Material should be available for at least three nights. Cold spells seldom last longer. The danger increases with the length of the cold spell. Trees and fruit become chilled and the last night of a cold spell is likely to be the one that does the damage. The freeze of December 23–27, 1906, lasted five nights, and in some sections the last night produced the lowest temperatures of the period. In some localities growers had exhausted their supply of firing material and, as a consequence, suffered a loss of fruit and young trees.

In cases of severe cold, fruit can not be saved, and trees can be preserved only by banking up the trunks by sand or soil. Small trees can be buried, and older trees can at least be protected above the parts where they have been budded.

The damage that will result from a given degree of cold varies with the condition and size of the tree. Well-grown trees in a dormant condition will stand without serious injury, except to foliage, a temperature of 18° to 20°, provided it is not too long continued. When the sap is flowing these temperatures are likely to kill growing trees, and are apt to cut small, unprotected trees to the ground. Well-grown trees in a dormant condition will stand without injury a temperature of 26° of continued cold, and can for several hours stand temperatures of 22° to 24°. When the sap is flowing these temperatures continued for several hours will be more or less injurious. Cold that will freeze fruit will injure small trees. In the freeze of December 23–27, 1906, temperatures in parts of the orange belt fell to 20°, and even to 18° and below, on one and in some cases two nights without injury to large trees, except to foliage, and with a surprisingly small loss of fruit and small trees. This freeze had been preceded by about two months of rainless weather, and found the trees in a dormant condition. The destructive freezes of preceding years usually followed periods of wet weather that promoted a flow of sap.

WORK IN MISSOURI.

In the Monthly Weather Review for May, 1910, Professor Howard, secretary of the Missouri State Board of Horticulture, expresses the following views relative to critical temperatures and the necessity of definite and detailed advance information in connection with frosts:

The horticulturists at the University of Missouri have closely determined the temperatures representing the danger point to fruit for the different stages in its development, as shown by the following brief summary of Circular No. 13, Missouri Experiment Station, 1909:

"Fully dormant peach buds can stand 8° or 9° below zero (F.). When they are appreciably swollen, zero is the danger point. When the buds are showing pink, they can stand 15° above zero. When the buds are almost open, 25° is the point of danger. When the petals are beginning to fall, 28° above zero is cold enough to cause uneasiness. When the petals are off, they can stand 30° above zero. When the 'shucks' (calyx tubes) are beginning to fall off, 32° above zero is the danger point."

It is estimated that the April, 1910, freeze caused a loss to the people of Missouri of $2,500,000, and it would mean thousands of dollars to the fruit growers in carrying out protective measures in the future to know just when to start the heaters in the orchards. It has been demonstrated that it is not a difficult matter, if one is prepared for it, and knows how to start the work, to raise the temperature in an orchard 5° to 6°, sufficient during almost any cold night in the spring to save many an orchard from injury. The question asked so often is, "When shall we start the heaters?" To start them too soon is expensive, but to start them too late would be more expen-

sive in the end. In this matter of advice the Weather Bureau must shoulder the burden, and it must take up the task with an earnestness that should prove successful. The forecaster must be thoroughly familiar, not only with the geographical location and topographical conditions of the section for which forecasts are made, but he must keep himself fully informed as to the progress of the fruit crop in its different stages of advancement. The state forecast issued by the district forecaster is all that is necessary for the public in general, but where specific information is desired it is not satisfactory. It will not suffice to simply send a weather message to a fruit locality in time of danger from low temperature, reading "Fair to-night and colder, with frost;" but the message must say *how cold* it is expected to be; for instance, something like the following: "Fair and colder to-night, with frost; temperature about 36°; no danger to fruit." And again, "Fair and colder to-night, with freezing temperature; 28° or lower by morning; start heaters after midnight."

While specifying the exact degree of cold expected may not be practicable for the district forecaster, owing to the extent of territory forecast for, and the limited time in which he has to make the forecast, it is being done, and can doubtless be brought to a high degree of accuracy, by the local forecasters.

DISTRIBUTION OF FORECASTS IN CALIFORNIA.

The system of frost warnings has reached a high degree of efficiency in California. The district forecaster sends telegrams each night to important fruit centers during the frost period, indicating the type of frost, whether light, heavy, or killing, the general weather conditions, and advice as to whether fire will be necessary.

To illustrate the way in which the information is distributed from the larger centers, the following statement is furnished by Mr. A. B. Wollaber, local forecaster at Los Angeles:

In California, where the changes in weather conditions can be only forecast for a short time in advance and where changes from one extreme to the opposite are often sudden, it has been found necessary to adopt a system whereby the local office can keep in touch at all times with the citrus industry. Here we find that the best means for accomplishing this purpose is the telephone, and in addition to the free distribution of our warnings by the long-distance companies operating in this section, we have an arrangement whereby we can reach the entire fruit belt within half an hour and disseminate our warnings through the medium of the fruit exchanges. The Weather Bureau pays for this service and therefore is in a position to demand and to receive prompt service. To accomplish this we have prepared a list of the exchanges in the fruit belt to which we send our warnings, and this list is on file with the chief operators of the long-distance companies. When we desire to send a warning we call for long-distance chief operator and give the message, which is immediately sent out. This system is used only in extreme cases, as when the regular maps and cards have been sent without a frost warning. That could occur in this manner: When a storm which gives indication of clouds or rain during the period covered by the morning forecast passes eastward more rapidly than was anticipated, causing a rapid clearing, and when later information from the district forecaster shows the probability of more severe conditions than anticipated on the morning reports.

Now, it happens in this section that not all ranches need protection at all times. The managers of the several fruit exchanges are personally familiar with conditions in their respective sections and are considered to be best qualified to disseminate this important information.

Special frost-warning cards are mailed to all post-offices, fruit exchanges, and railroad depots before the frost period begins. There has grown up a personal relationship between the growers and the local office, and through this much good has been accomplished.

MISCELLANEOUS.

A TWO YEARS' STUDY OF SPRING FROSTS AT WILLIAMSTOWN, MASS.

A valuable paper by Prof. W. I. Milham on the above subject was published in the Monthly Weather Review for August, 1908. The special purpose of the investigation was the critical study of the spring frosts, first, to test the prevalent belief that the probable minimum temperature could be computed from the maximum and dew-point, or from the reading of the wet-bulb thermometer of the afternoon before the frost; second, to study the variation in the severity of frost at different places within a small area; and, third, to study the variation in the severity of frost with varying distances above the ground. Professor Milham came to the following conclusions:

(1) The cool nights of spring when a frost might be expected are very dry, and the dew-point lies so low that it plays practically no part in determining the minimum temperature.

(2) The amount of the drop from the maximum to the following minimum is very far from a constant, even if the characteristics of these nights seem very nearly the same. In each case in estimating the probable drop one must take into account the probable amount of cloud, the probable wind velocity, the possibility of a change in wind direction, and possibly other things.

Milham holds that two processes operate to produce the cooling, first, the importation of cooler air, and, second, the radiation of heat from the ground and the cooling of the air next to it by conduction. It might seem that the cool nights during which frosts seem probable could be divided into two groups, depending upon whether the importation of cold air or radiation from the ground was playing the larger part in producing the cooling. It is found, however, that both processes are nearly always operative. It may not be out of place at this point to call attention to the fact that the heat which has been absorbed by the ground is re-emitted as radiation of long wave lengths, from 4 μ to 20 μ.

Milham calls attention to the fact that the temperature of vegetation in the open, near the ground, in the coldest part of the tract may be 11° lower than the estimated minimum in a given shelter.

An interesting paper upon temperature variations near the surface of the soil during the formation of frost, by Mr. Dewey A. Seeley, local forecaster, can be found in the Monthly Weather Review for August, 1908. He is of the opinion that a considerable amount of

the heat given off, during the condensation of vapor as frost and dew, is retained in saturated air near the ground level and also by the foliage of the plants themselves. The influence of a surface covering of vegetation on the temperature of the soil was marked. It was found that the soil beneath the grass was warmer both at night and during the day than the naked soil; but the temperature of the air above the grass was lower in both cases than over bare ground.

The dew-point as recorded at 7 p. m. and the minimum temperature registered the following morning for sixty days were tabulated and the differences computed. From these it appears that on the average the minimum temperature went 2.9° below the 7 p. m. dew-point. In but three cases did the temperature fall more than 10° below the dew-point recorded at 7 p. m. The thermograph traces made during clear weather with light wind show that the temperature fall goes on much more rapidly during the first part of the night, before the dew-point is reached.

[A list giving the titles of all Farmers' Bulletins available for distribution will be sent free upon application to a Member of Congress or the Secretary of Agriculture.]

104

O

R

Issued November 17, 1911

Reserve

U. S. DEPARTMENT OF AGRICULTURE.

FARMERS' BULLETIN 104.

NOTES ON FROST.

LIBRARY
RECEIVED
JAN 21 1920
U. S. Department of Agriculture

BY

E. B. GARRIOTT,
Professor of Meteorology, Weather Bureau.

Revised August, 1911.

By ALEXANDER G. McADIE,
Professor of Meteorology, Weather Bureau.

WASHINGTON:
GOVERNMENT PRINTING OFFICE.
1911.

LETTER OF TRANSMITTAL.

U. S. DEPARTMENT OF AGRICULTURE,
OFFICE OF THE WEATHER BUREAU,
Washington, D. C., August 22, 1911.

SIR: I have the honor to recommend that the manuscript herewith be published as a revised edition of Farmers' Bulletin No. 104.

In compliance with directions contained in your letter of March 29, 1899, a paper on the subject of frost, prepared by Prof. E. B. Garriott, Weather Bureau, was submitted and published as Farmers' Bulletin No. 104. This bulletin furnished valuable information regarding methods of protection from frost. Methods advocated in previous publications were mainly the results of experiments conducted in California orchards. Bulletin No. 104 was designed to be of value in all sections of the country rather than in restricted districts.

As the California experiments quoted were perhaps the most thorough ever made in this country, a description of the methods employed and the results obtained is reprinted in brief in this paper, together with results of similar experiments made in the citrus-fruit and truck-growing districts of the Gulf States, where varying climatic conditions have been found to produce somewhat different results. Additional information personally acquired by Prof. Garriott during a visit to Florida in the winter of 1906–7, and certain climatic data, also form features of this paper in its revised form.

The development of orchard heaters and other devices now successfully used in saving crops from injury by frost necessitates a new issue of this bulletin. Owing to the death of Prof. Garriott, the revision has been undertaken at my direction by Prof. McAdie.

Your wishes that the subject be treated in a popular way and divested of all technicalities, and made applicable to all portions of the United States, have been complied with so far as the nature of the subject would permit.

Respectfully,

WILLIS L. MOORE,
Chief, Weather Bureau.

Hon. JAMES WILSON,
Secretary of Agriculture.

CONTENTS.

	Page.
How frost is formed	5
Seasons of frost	7
When to expect frost	10
Influence of soil and vegetation on minimum temperature	14
Dates of early autumn and late spring frosts	15
Protection from frost	17
Devices for preventing rapid radiation of heat	17
Glass screens	18
Cloth screens	18
Lath screens	18
Paper cover	18
Other methods	19
Devices for adding heat to the air	19
Antifrost candle	24
Devices for adding moisture to the air	25
Portable smudge fires	25
General observations and practical results	26
Work in Florida	27
Work in Missouri	29
Distribution of forecasts in California	30
Orchard heating in Indiana	31
Work in Oregon	32
Work in Utah	33
Variability of frost injury on fruit buds	33
Miscellaneous	34
A two years' study of spring frosts at Williamstown, Mass	34

ILLUSTRATIONS.

		Page.
FIG.	1. Chart showing average date of first killing frost in autumn	8
	2. Chart showing average date of last killing frost in spring	9
	3. Diagram of Kunkel orchard, Wichita, Kans	22
	4. Diagram of plat of Thomas Orchard Co., Wichita, Kans	23

NOTES ON FROST.

HOW FROST IS FORMED.

The atmosphere of the earth always contains more or less moisture in an invisible form. When at a considerable elevation above the earth this moisture, or aqueous vapor, is condensed, clouds are formed; when the process of condensation is more active and the temperature of the air is above freezing, rain falls; and when the temperature of the air is below freezing, snow is produced. When the moisture of the air in immediate contact with the earth is condensed at temperatures above freezing, dew is formed; when at temperatures below freezing, frost is deposited. Frost is, therefore, the moisture of the air condensed at freezing temperatures upon plants and other objects near the surface of the earth.

In the process of frost formation the temperature of the air a few feet above the earth is commonly several degrees above freezing. The surfaces upon which frost is deposited must, however, possess freezing temperatures. The manner in which frost is deposited on plants and other objects is very similar to that in which the air moisture of a room is frozen and deposited upon window glass, the temperature of which has been reduced to freezing by the out-of-doors cold. In the case of the frost on the window glass the process is one that can readily be understood. Some explanation, however, is required of the formation of frost, which requires freezing temperature, when at times a temperature above freezing is registered a few feet above the surfaces upon which frost appears.

There are several processes by which the temperature of plants may be reduced below the temperature of the air which surrounds them. The most important of these processes is radiation, by means of which heat escapes from objects and passes into the surrounding air.

In the frost-forming process heat from the sun which is absorbed by the earth and by plants during the day is lost by radiation at night. During the day the earth both absorbs and reflects the heat received from the rays of the sun, and the lower stratum of air is warmed by this reflected heat. During the night, when no direct heat is received from the sun's rays, the lower air stratum receives no

reflected heat, and at the same time heat which has been absorbed by the earth is radiated, or, in other words, it rises through the overlying air, though not necessarily and entirely as sensible heat; that is, heat perceptible to the senses. In the frost-forming process there is another very important factor in the production of cold; that is, the evaporation of moisture from the earth and from plants. The nature and composition of frost require that the air in immediate contact with the surfaces upon which frost forms shall contain moisture. This is moisture that has been stored in the earth in visible form, i. e., in the form of water, and which has escaped from the earth, not as visible moisture, but by the process of evaporation and in the form of aqueous vapor. Some part of the heat stored by the earth during the day is, therefore, lost in evaporating the moisture contained in plants and in the earth. The heat expended in this process is not sensible, but latent. Heat is rendered latent when it is used to boil away water, and it becomes sensible or liberated heat when the water vapor again condenses into water. It appears, therefore, that the temperature of surfaces upon which frost forms, and of the air in immediate contact with them, is lowered by the evaporation of moisture from the surfaces. The influence of the process does not extend to air a few feet above the ground.

Another method by which plants lose their heat is by convection, by means of which they are chilled by contact with colder air. This process, while important in the presence of freezing air temperatures, can scarcely be considered a factor in the formation of frost proper, which is usually accomplished when temperature observations show the air to be above freezing.

Air arranges itself according to its density or weight. Air in immediate contact with the earth becomes heavier as its temperature is lowered by the radiation of heat and settles in depressions or valleys and over lowlands, causing, at times, frost in low-lying districts, while neighboring higher grounds escape the visitations.

NOTE.—A given quantity of heat absorbed by several substances of different specific heat will produce a different temperature in each; the lower the specific heat of the substance the higher will be its temperature; and besides, different substances have different coefficients of asorption and of reflection. Therefore under the same insolation rocks, clay, vegetation, and other substances come to temperatures that often differ by many degrees.

The air next to earth largely partakes of the temperature of the surface, and in consequence, when the sky is clear, so that incoming and outgoing radiations may progress freely, and the air is but little disturbed by wind, wide variations in the temperature of the thin stratum of lower air may occur over adjacent plots of ground of precisely the same elevation but different covering.—Moore's Descriptive Meteorology, p. 108.

SEASONS OF FROST.

Generally speaking, agricultural products in some part of the United States are menaced by frost from the time nature begins to extend her mantle of green from the citrus regions of the South toward the grain fields of the North until the verdured area contracts southward before the first winds of winter; from the hour that seeds germinate and buds blossom until crops are gathered; from early spring until late fall. As a rule, the critical periods are during the early growth of plants and when the fruit is in the bloom, and again when crops are maturing. In the first instance vegetation is threatened by the frosts of early spring, and in the latter by the frosts of autumn.

Considering the several agricultural sections in greater detail we find that damaging frost is likely to occur in the north half of the Florida Peninsula, the region immediately bordering the coast of the Gulf of Mexico, and in the interior of California from the latter part of October until the early part of April, and in the North Pacific Coast States as early as the middle of October and as late as the last week of April. From December until early in March the frost limit and even the line of freezing temperature may, at long intervals, be extended southward over central and southern Florida. The period of damaging frosts in the interior of the South Atlantic and Gulf States extends from November to April. From October to April the region in which agricultural products are subject to damage by frost is extended to the southern lake region, over the Upper Mississippi and Lower Missouri Valleys, Kansas, and Nebraska, and from September to May frost visitations are usually confined to the extreme Upper Mississippi, Upper Missouri, and Red River of the North Valleys, and the Rocky Mountain and plateau regions from central New Mexico and the Texas Panhandle northward. (See figs. 1 and 2.)

The occurrence of summer frosts in mountain regions and in extreme northern districts of the country is common. The following are notable instances of summer frosts in agricultural districts:

August 22 to 24, 1890, a frost-bearing cool wave advanced from the Dakotas over the lake region and the interior, or mountain districts, of the middle Atlantic States.

August 28 to 31, 1892, a cool wave advanced from the Rocky Mountain and plateau regions, with frost from New Mexico to the Dakotas, Iowa, Minnesota, northern Wisconsin, and northern Michigan.

August 8, 1904, frost damaged vegetation in low-lying lands in the States of the upper lake region.

Bulletin P, of the Weather Bureau, gives a chronological statement of the more important cold periods that have been experienced in the United States, and contains charts that present conditions that preceded and attended the principal cold waves and frosts of the 15 years from 1888 to 1902.



FIG. 1.—Chart showing average date of first killing frost in autumn.



FIG. 2.—Chart showing average date of last killing frost in spring.

Bulletin V, of the Weather Bureau, issued in 1911, gives in more detail frost data and length of the crop-growing season as determined from the average of the latest and earliest dates of killing frost.

WHEN TO EXPECT FROST.

As the conditions which produce damaging frost are subject to modifications which are as numerous as the kinds of crops raised, and as varied as local topography, local climate, and local soil conditions in the various sections of the United States, the writer in attempting to treat briefly a subject which is so diversified in its aspects is confronted with many difficulties. Suffice it to say that with other atmospheric conditions favorable for its occurrence frost may, as a rule, be expected when temperature, as reported by stations of the Weather Bureau, falls to a point 8 to 10° above the freezing point. As previously stated, the surfaces upon which frost is deposited must be at or below freezing temperature, although the temperature of the air a few feet above may be several degrees above freezing. It is the temperature of the air, in some instances many feet above the ground, that is given by the Weather Bureau observations. Another atmospheric condition favorable for the occurrence of frost is a clear, cloudless, and comparatively calm night. Clouds retard radiation or loss of heat from plants and act as a screen in preventing the heat collected from the sun's rays during the day from escaping into the upper air. When clouds are not present and a withdrawal of the sun's rays causes a rapid cooling of the air at moderate elevations, the warmer air which collects near the surface of the earth during the day rises, and the cooler upper air, owing to its greater density or weight, settles to the earth. It will be noted that clouds not only prevent the escape of the warm air in immediate contact with the earth, but also blanket, as it were, the upper part of the lower air stratum. Calm or comparatively still air is a condition which favors the formation of frost. On windy nights the air is disturbed and is not permitted to arrange itself in layers according to its density, with the densest and coldest air near the surface of the earth; it is kept mixed up by the wind.

A very interesting fact, which illustrates the manner in which cold air settles by its greater weight to the surface of the earth and allows the warmer air to assume higher levels, causing frost in low grounds and granting to the higher grounds a comparative immunity from frost, is set forth in the following article, by Prof. Cleveland Abbe, in the Monthly Weather Review for December, 1893, entitled "Thermal belts, frostless belts, or verdant zones":

These are local names given to certain regions on mountain sides within which nocturnal frost rarely or never occurs in the springtime, although freezing temperatures occur in the wintertime. Consequently tender vegetation flourishes

with remarkable vigor in these regions. The following are the only references to thermal belts in the United States at present known to the editor, but as such regions are specially important to the horticulturist and agriculturist it is hoped that the correspondents of the Weather Bureau will bring these thermal belts to notice wherever they occur in order that their meteorological peculiarities may be better understood.

In the Agricultural Report of the Patent Office for 1861 Mr. Silas McDowell, of Franklin, Macon County, N. C., describes the verdant zone in that county in the valley of the Little Tennessee River. It occupies the region between 300 and 700 feet above the valley of the river, which latter is about 2,000 feet above sea level. On tracing this zone up among the smaller tributaries of the Tennessee River he found that in the higher valleys, where the bottom land is about 3,900 feet above sea level, the verdant zone lies between 4,000 and 4,100 feet. Within this zone frost never injures the vegetation, and the most tender grapes never fail to produce abundant crops.

Prof. J. W. Chickering, jr., in the bulletin of the Philosophical Society of Washington, March, 1883, and in the American Meteorological Journal, volume 1, describes the following thermal belt: "In Polk County, N. C., along the eastern slope of the Tryon Mountain Range, in latitude north 35, the thermal belt begins at the base of the mountain, at an elevation of 1,200 feet above the sea, and extends up 2,200 feet, being most perfect at about 1,500 feet. It is about 8 miles long, and is distinguished by magnificent flora, such as would be characteristic of a point 3° south of the actual latitude."

Prof. John Leconte, of Berkeley, Cal., in Science, volume 1, page 278, states that at Flat Rock, near Hendersonville, Henderson County, N. C., on the flank of the mountain spur adjacent to the valleys of the Blue Ridge, he also observed a frostless zone. The valley is about 2,200 feet above sea level, and the thermal belt is 200 to 300 feet above the valley.

In the American Meteorological Journal, volume 1, Mr. S. Alexander describes a thermal belt in which the peach tree flourishes in the southeastern portion of Michigan. He shows that the cold island discovered by Winchell in that region is really the bottom of a topographical depression into which the cold air settles. It is a long valley surrounded by a belt of elevated country from 50 to 600 feet above Lakes Michigan and Huron. The valley and the isotherms trend northeast and southwest from Huron County through Sanilac, Lapeer, Oakland, Livingston, and Washtenaw to Hillsdale County. The highlands of this region are all much freer from frost than the lowlands, and all much more favorable for early vegetation. He does not state that any point is high enough to be above the thermal belt, but that, in general, two equal parallel thermal belts inclose the cold island between them.

The thermal belts of California are described in various publications of the Weather Bureau, particularly Bulletin L, "Climatology of California."

In the various numbers of the Monthly Weather Review and in the various summaries published by section directors and district editors special attention has been given to zones of little injury from frost. In nearly every section there are some specially favored localities. One example will serve to illustrate. In the Monthly Weather Review for May, 1910, Prof. Howard, secretary of the Missouri State Board of Horticulture, calls attention to the fact that certain sections in that State, for no apparent reasons based upon

topography or isotherms, nearly always escape injury. He instances particularly the famous peach region about the town of Koshkonong, in Oregon County. Mr. George Reeder, section director, after a careful examination of climatic data, comes to the conclusion that the apparent immunity of orchards in the neighborhood under discussion from damaging temperatures in the spring is probably due to local air drainage, the cold air being drained away from the orchards. He gives tables showing that the cold air spreads out and is drained away down the southern slope of the Ozark Plateau, sometimes more toward the southwestward, then toward the south, and again the flow may affect only the southeast. Mr. Reeder states:

It appears as if the cold air currents had divided, one part rolling toward the White River Valley, as indicated by the temperature at Hollister, and the other part going toward the Black River Valley, as indicated by the temperature at Doniphan. Instead of the favorable peach belt, if any, extending east and west, as suggested by Prof. Howard, it probably extends northerly and southerly, or, in other words, it lies along the Ozark border, a narrow strip of tableland having elevations ranging between 1,000 feet, on which are situated Koshkonong, Mo., and Mammoth Springs, Hardy, and La Crosse, Ark. This strip bends not far from La Crosse, and thence extends northwestward above the White River Valley. The temperature on this border does not fall so low at times as does that at lower as well as higher altitudes.

The cold of April 25 and 26, 1910, that caused so much damage to the fruit crop of Missouri was quite uniformly distributed over the section under discussion, as may be seen from the following table:

Counties.	Stations.	Temperature.	Date.
		° F.	
McDonald	Dean	28	Apr. 26
Taney	Hollister	31	Do.
Howell	Olden	27	Apr. 25
Oregon	Koshkonong	29	Do.
Ripley	Doniphan	32	Do.

The cold spell was accompanied by cloudy skies and rain mixed with snow, and the movement of the cold currents of air was quite different from that which takes place under a clear sky, or when there is no rain or snow. It will be observed that during cold periods, with more or less cloudiness and moisture, the temperature on the hills may fall to a lower value than the temperature in the valleys; on the other hand, had clear weather prevailed during the nights of April 25 and 26, the temperature conditions would have been reversed, the valleys probably being much colder. It is also quite clear, had the fruit in the valleys escaped winter injury it would have received no more damage from the freeze of April than was sustained in the neighborhood of Koshkonong.

Prof. Abbe states in Monthly Weather Review for December, 1903:

It is generally conceded that these thermal belts depend both upon the drainage of cold air downward into the lower valleys and the freedom of radiation from the surface of the ground to the clear sky overhead. During a still night, when frosts occur, the surface of the hillside cools by radiation, and hence cools the air in contact with it; the latter flows downward as long as its cooling by

radiation and conduction exceeds its warming by compression. Inasmuch as its cooling depends on contact with a still colder soil or plant, it soon accumulates in the lowlands as a layer of cold air, which grows thicker during the night by the steady addition of the thin layer of descending air in contact with the ground on the hillsides. The warmer air, which has not yet had an opportunity to cool by contact with the ground, floats on top of the cold mass; it spreads out toward the hills, and is continuously furnishing its heat to the adjacent hillsides as fast as it comes in contact with them before it also cools and descends. The formation of the thermal belt seems to depend largely upon this gentle circulation during the nighttime. The lower limit of the belt is defined by the depth of the accumulation of cold air in the confined valley and rises higher in proportion as the night is clearer and longer, and also in proportion as the valley is more or less perfectly inclosed. The upper limit of the thermal belt may depend upon the strength of the wind and the general temperature of the air. But if there be no wind, then it depends equally on the freedom of radiation to the clear sky and on the above-described circulation of air.

The facts quoted above show that frost may be expected on low grounds at times when higher grounds escape the visitations, and the lesson they teach is that early and tender crops should, so far as may be practicable, be confined to crests, hillsides, and mountain sides, and later and hardier crops to the lowlands and valleys.

Local climate, as it is influenced and regulated by the proximity of bodies of water, must be given great weight in calculations regarding frost. Frost is less likely to occur in localities swept by moisture-laden air which has crossed a considerable body of water. This is more especially the case in the fall of the year, when the temperature of bodies of water is reduced very slowly; and in the South, where the water temperature continues relative high throughout the year, the influence of the water is especially marked. During the colder months air is not only warmed in crossing considerable bodies of water, but it also absorbs moisture, which, although invisible in the form of water vapor, has the effect of retarding the radiation of heat from the earth. In fact, so pronounced is the influence of water and water vapor that localities which with reference to exposure to west and northwest winds are, as it were, in the lee of large bodies of water are comparatively free from frost visitations. On the keys of the east Florida coast, which are protected on the west by lagoons, rivers, lakes or bays, frost is said to be unknown, and throughout the central and northern sections of the United States, localities which are protected on the west and northwest by bodies of water of considerable size, are not only favored, by reason of their position, with conditions which do not promote the formation of frost, but also receive the benefits derived from heavier falls of snow.

Local soil conditions constitute, to a certain extent, a factor in the formation of frost. As a rapid loss of heat is promoted by an active evaporation of moisture, it is evident that, with other conditions

equal, frost is more likely to occur on damp than on dry ground, provided, however, that the ground is not too moist, for in the latter event the amount of moisture evaporated and added to the air would have a tendency to retard the radiation of heat from the earth. And herein lies the distinction, so far as frost formation is concerned, between moist air and moist soil. An excess of moisture in the air, in preventing, to a degree, the radiation of heat, is unfavorable to the formation of frost. As frost is the moisture of the air in immediate contact with the earth condensed at freezing temperature, it follows that the earth, from which the moisture of the air is drawn, must contain more or less water, and it is evident that damaging frost will occur with a limited rather than with a large quantity of moisture arising from the earth and a moderately moist and comparatively still air. Both of these conditions usually obtain following, but not too closely following, the rains of spring and autumn. The character of the soil as regards its capacity for retaining moisture must also be taken into account, more especially during periods of comparatively dry weather. Moist soil, as well as the vegetation which springs from it, is chilled to a lower temperature by the evaporation of moisture, and is therefore more subject to visitations of frost than soil which is dry, or into which water penetrates deeply, leaving the surface dry, or soil which sheds the rain and does not absorb and hold water. It is important to note, however, that very moist soil, or soil which contains a large amount of surface water, is, owing to the quantity of moisture yielded to the air, not so subject to the heavy and damaging frosts which visit plants on moderately damp ground or ground that has a small capacity for moisture.

INFLUENCE OF SOIL AND VEGETATION ON MINIMUM TEMPERATURE.

Prof. W. L. Moore, while engaged in studying frost formation in the cranberry bogs of Wisconsin, in 1891 to 1894, came to the conclusion that for a given area the occurrence of frost depended more upon the character of the surface, its covering, and the degree of heat and moisture to which it had been subjected for several days preceding than to the general temperature and pressure of the air, one field receiving an injurious frost, another a light frost, and still another none at all, while apparently the same weather conditions prevailed at a moderate distance above the ground. Experiments carried on since 1906 by Prof. Cox in connection with the protection of cranberries show that in the bogs there may be surface differences of from 5° to 10° in temperature, while at an elevation of 3 feet these differences disappear. Prof. Cox is of opinion that variation in temperature is due largely to differences in temperature of the soil

or covering. It is as if heaters of varying power were scattered over the bog, giving off heat to the air immediately above. Investigations carried on at Berlin, Cranmoor, and Mather show that in sections which were carefully weeded and cleaned and were free from moss the temperatures were higher. Progressive cranberry growers find that by cultivating, draining, and sanding, low night temperatures are prevented.

Bulletin T, of the Weather Bureau, issued in 1910, on Frost and Temperature Conditions in the Cranberry Marshes of Wisconsin, gives the results of an investigation covering several years' field work.

DATES OF EARLY AUTUMN AND LATE SPRING FROSTS.

The following table shows for stations of the Weather Bureau throughout the United States dates of first killing frost in autumn and of last killing frost in spring; also the average dates of first killing frost in autumn and of last killing frost in spring:

Dates of first killing frost in autumn and of last killing frost in spring at Weather Bureau stations throughout the United States.

Stations.	Date of earliest recorded killing frost in autumn.	Date of latest recorded killing frost in spring.	Average date of first killing frost in autumn.	Average date of last killing frost in spring.
Eastport, Me	Sept. 8	June 19	Oct. 12	Apr. 28
Northfield, Vt	Aug. 27	June 7	Sept. 16	May 13
Portland, Me	Sept. 11	May 31	Oct. 18	Apr. 14
Boston, Mass	Sept. 30	May 11	Oct. 22	Apr. 26
Nantucket, Mass	Oct. 1	Apr. 24	Nov. 5	Apr. 10
Block Island, R. I	Oct. 30	May 11	Nov. 16	Apr. 12
Narragansett, R. I	Sept. 21	May 30	Oct. 16	Apr. 23
New Haven, Conn	Sept. 15	...do	Oct. 17	Apr. 20
Binghamton, N. Y	Sept. 16	May 21	Oct. 6	Apr. 27
Albany, N. Y	Sept. 23	May 30	Oct. 17	Apr. 24
New York, N. Y	Oct. 5	Apr. 30	Nov. 6	Apr. 10
Harrisburg, Pa	Oct. 3	Apr. 26	Oct. 24	Do.
Philadelphia, Pa	...do	Apr. 29	Oct. 30	Apr. 8
Atlantic City, N. J	Oct. 1	Apr. 25	Nov. 4	Apr. 11
Baltimore, Md	Oct. 6	May 3	...do	Apr. 4
Washington, D. C	Oct. 2	May 11	Oct. 21	Apr. 7
Lynchburg, Va	Oct. 4	May 7	Nov. 1	Apr. 14
Norfolk, Va	Oct. 15	Apr. 26	Nov. 12	Mar. 27
Wytheville, Va	Sept. 14	May 26	Oct. 7	Apr. 23
Charlotte, N. C	Oct. 8	Apr. 26	Nov. 4	Apr. 1
Raleigh, N. C	...do	May 6	...do	Apr. 5
Hatteras, N. C	Nov. 7	Apr. 19	Dec. 11	Feb. 28
Wilmington, N. C	Oct. 16	May 1	Nov. 15	Mar. 27
Charleston, S. C	Nov. 9	Apr. 2	Nov. 30	Mar. 1
Columbia, S. C	Oct. 19	Apr. 17	Nov. 8	Mar. 23
Augusta, Ga	Oct. 8	...do	Nov. 9	Mar. 20
Savannah, Ga	Nov. 1	Apr. 5	Nov. 26	Feb. 27
Jacksonville, Fla	Nov. 12	Apr. 6	Dec. 6	Feb. 14
Jupiter, Fla	Nov. 18	Apr. 7	Dec. 29	Do.
Atlanta, Ga	Oct. 11	Apr. 15	Nov. 7	Mar. 24
Macon, Ga	Nov. 5	Apr. 18	Nov. 14	Mar. 15
Tampa, Fla	Nov. 28	Mar. 19	Jan. 9	Feb. 8
Pensacola, Fla	Nov. 12	Apr. 6	Dec. 5	Feb. 23
Mobile, Ala	Oct. 31	Mar. 28	Nov. 30	Feb. 24
Montgomery, Ala	Oct. 21	Apr. 5	Nov. 8	Mar. 11
Meridian, Miss	Oct. 8	Apr. 10	Nov. 2	Mar. 20
Vicksburg, Miss	Oct. 19	Apr. 6	Nov. 12	Mar. 6
New Orleans, La	Nov. 19	Mar. 27	Dec. 15	Jan. 24
Shreveport, La	Oct. 20	Apr. 2	Nov. 11	Mar. 4
Fort Smith, Ark	Oct. 15	Apr. 6	Nov. 4	Do.
Little Rock, Ark	Oct. 22	Apr. 14	Nov. 10	Mar. 20

Dates of first killing frost in autumn and of last killing frost in spring at Weather Bureau stations throughout the United States—Continued.

Stations.	Date of earliest recorded killing frost in autumn.	Date of latest recorded killing frost in spring.	Average date of first killing frost in autumn.	Average date of last killing frost in spring.
Palestine, Tex	Oct. 20	May 30	Nov. 13	Mar. 13
Galveston, Tex	Dec. 4	Mar. 1	Dec. 25	Feb. 5
San Antonio, Tex	Nov. 9	Mar. 20	Nov. 30	Feb. 25
Corpus Christi, Tex	Nov. 30	Mar. 19	Dec. 25	Feb. 27
Memphis, Tenn	Oct. 2	Apr. 16	Oct. 28	Mar. 24
Nashville, Tenn	Oct. 8	May 14	Oct. 24	Apr. 2
Chattanooga, Tenn	Sept. 30	...do....	Oct. 26	Do.
Knoxville, Tenn	Oct. 1	Apr. 24	Oct. 27	Apr. 3
Louisville, Ky	Sept. 24	May 14	Oct. 29	Apr. 6
Lexington, Ky	Sept. 30	May 20	Oct. 23	Apr. 18
Evansville, Ind	...do....	Apr. 21	Oct. 30	Apr. 10
Indianapolis, Ind	Sept. 21	May 21	Oct. 19	Apr. 16
Cincinnati, Ohio	Sept. 30	Apr. 24	Oct. 25	Apr. 14
Columbus, Ohio	Sept. 21	May 17	Oct. 16	Apr. 16
Elkins, W. Va	Sept. 28	May 24	Oct. 10	May 18
Parkersburg, W. Va	Sept. 24	May 22	Oct. 17	Apr. 11
Pittsburgh, Pa	Sept. 25	May 29	Oct. 19	Apr. 26
Oswego, N. Y	...do....	...do....	...do....	Apr. 27
Rochester, N. Y	Sept. 26	May 30	...do....	May 1
Buffalo, N. Y	Sept. 23	May 29	Oct. 16	Apr. 25
Erie, Pa	Oct. 12	May 17	Oct. 29	Apr. 22
Cleveland, Ohio	Oct. 2	May 22	Oct. 31	Apr. 16
Sandusky, Ohio	Oct. 8	May 17	Oct. 26	Apr. 14
Toledo, Ohio	Sept. 9	May 29	Oct. 15	Apr. 24
Detroit, Mich	Sept. 21	May 31	Oct. 11	Apr. 30
Port Huron, Mich	Sept. 22	June 6	Oct. 9	May 8
Alpena, Mich	Sept. 6	June 9	Sept. 26	May 14
Sault Ste. Marie, Mich	Sept. 5	May 29	Sept. 24	May 16
Marquette, Mich	Aug. 22	June 11	Oct. 2	May 15
Escanaba, Mich	Sept. 9	June 16	Oct. 1	May 14
Green Bay, Wis	Sept. 16	May 30	Oct. 4	May 5
Grand Haven, Mich	Sept. 23	May 28	Oct. 10	Apr. 28
Milwaukee, Wis	Sept. 25	May 29	...do....	Apr. 29
Chicago, Ill	Sept. 18	...do....	Oct. 15	Apr. 18
Duluth, Minn	Sept. 15	June 8	Oct. 5	May 3
Minneapolis, Minn	Sept. 13	May 20	Oct. 8	Apr. 27
St. Paul, Minn	Sept. 20	May 25	Oct. 5	May 6
Madison, Wis	Sept. 29	May 13	Oct. 17	Apr. 21
La Crosse, Wis	Sept. 21	June 1	Oct. 8	May 2
Dubuque, Iowa	Sept. 27	May 21	Oct. 13	Apr. 20
Davenport, Iowa	Sept. 18	May 22	Oct. 14	Apr. 22
Des Moines, Iowa	Sept. 12	...do	Oct. 8	Apr. 28
Keokuk, Iowa	Sept. 18	May 4	Oct. 15	Apr. 13
Hannibal, Mo	Sept. 30	May 14	...do....	Apr. 14
Springfield, Ill	Sept. 25	May 22	Oct. 16	Apr. 20
St. Louis, Mo	Sept. 30	...do....	Oct. 27	Apr. 2
Cairo, Ill	...do....	Apr. 19	Oct. 28	Mar. 29
Springfield, Mo	...do....	May 19	Oct. 13	Apr. 16
Columbia, Mo	Sept. 28	May 9	Oct. 15	Apr. 14
Kansas City, Mo	Sept. 30	May 4	Oct. 24	Apr. 10
Topeka, Kans	Sept. 28	May 19	Oct. 13	Apr. 8
Wichita, Kans	Sept. 23	May 15	Oct. 19	Apr. 6
Concordia, Kans	Sept. 27	May 19	Oct. 14	Apr. 24
Omaha, Nebr	Sept. 18	...do....	Oct. 12	Apr. 15
Lincoln, Nebr	Sept. 12	May 10	Oct. 9	Apr. 18
Valentine, Nebr	...do....	June 21	Sept. 18	May 9
Sioux City, Iowa	Sept. 13	May 21	Sept. 27	May 1
Huron, S. Dak	Aug. 23	June 8	Sept. 20	May 13
Pierre, S. Dak	Sept. 12	May 19	Sept. 30	Apr. 30
Moorhead, Minn	Aug. 25	June 8	Sept. 22	May 14
Bismarck, N. Dak	Aug. 23	June 7	Sept. 21	May 15
Williston, N. Dak	Aug. 18	June 10	Sept. 14	May 18
Havre, Mont	Aug. 27	June 6	Sept. 13	May 14
Helena, Mont	Sept. 5	June 9	Sept. 25	May 10
Rapid City, S. Dak	Sept. 13	May 21	Sept. 20	May 1
Lander, Wyo	Aug. 23	June 18	Sept. 11	May 19
Salt Lake City, Utah	Sept. 22	...do....	Oct. 18	Apr. 23
Grand Junction, Colo	Sept. 27	May 15	Oct. 21	Apr. 14
Cheyenne, Wyo	Aug. 29	June 11	Sept. 16	May 22
North Platte, Nebr	Sept. 10	May 23	Sept. 28	May 1
Denver, Colo	Sept. 12	June 6	Oct. 5	May 6
Amarillo, Tex	Oct. 16	May 23	Nov. 1	Apr. 16
Pueblo, Colo	Sept. 12	...do....	Oct. 5	Apr. 26
Dodge, Kans	Sept. 23	May 27	Oct. 15	Apr. 17

Dates of first killing frost in autumn and of last killing frost in spring at Weather Bureau stations throughout the United States—Continued.

Stations.	Date of earliest recorded killing frost in autumn.	Date of latest recorded killing frost in spring.	Average date of first killing frost in autumn.	Average date of last killing frost in spring.
Oklahoma, Okla.	Oct. 7	Apr. 30	Oct. 31	Apr. 2
Fort Worth, Tex.	Oct. 22	May 1	Nov. 19	Mar. 18
Abilene, Tex.	Oct. 24	Apr. 16	Nov. 15	Mar. 15
El Paso, Tex.	Oct. 23	Apr. 22	Nov. 10	Mar. 20
Santa Fe, N. Mex.	Sept. 27	May 18	Oct. 19	Apr. 15
Yuma, Ariz.	None.	None.	None.	None.
Phoenix, Ariz.	Nov. 9	Mar. 31	Dec. 7	Feb. 18
Spokane, Wash.	Sept. 7	June 8	Oct. 17	Mar. 23
Walla Walla, Wash.	Sept. 28	May 12	Nov. 7	Apr. 4
Seattle, Wash.	Oct. 18	May 27	Nov. 22	Mar. 21
Portland, Oreg.	Oct. 13	May 9	Nov. 16	Mar. 17
Tatoosh Island, Wash.	Nov. 1	Apr. 19	Dec. 15	Mar. 14
Roseburg, Oreg.	Sept. 25	May 16	Oct. 30	Apr. 15
Baker City, Oreg.	Sept. 4	June 24	Sept. 27	May 25
Winnemucca, Nev.	Aug. 22	June 20	Sept. 24	May 15
Eureka, Cal.	Nov. 7	May 1	Dec. 8	Apr. 2
Red Bluff, Cal.	Nov. 15	Apr. 25	Dec. 6	Feb. 26
Sacramento, Cal.	Oct. 28	Apr. 26	Nov. 15	Feb. 16
San Francisco, Cal.	Nov. 18	Apr. 20	Dec. 26	Jan. 8
Fresno, Cal.	...do....	Apr. 14	Dec. 15	Mar. 4
San Luis Obispo, Cal.	Nov. 19	Apr. 8	Dec. 27	Feb. 20
Los Angeles, Cal.	None.	...do....	None.	Apr. 8
San Diego, Cal.	None.	None.	None.	None.

PROTECTION FROM FROST.

A discussion of methods of protection from frost calls for a consideration in great detail, not only of the several approved processes of protection and the extent to which they can be applied in connection with the various crops, but also of the value of these processes under the varying conditions of local topography, local climate, and local soil conditions.

It is evident, in the light of the foregoing statements, that fruit and vegetable growers in rolling and hilly country, or when located in the lee, as regards west to northwest winds, of bodies of water, can, to some extent, so place in their earlier and tenderer fruits and crops that they will, in the first instance, avoid the lowlands and valleys, and in the second take advantage of the more moderate and moist air that crosses water surfaces. For further protection the grower must depend upon artificial appliances. These appliances are designed to produce the following effects or results: To prevent a rapid radiation of heat from the earth; to charge the air with moisture; to warm the air; to create artificial drafts, whereby the air is mixed and the cold air is not allowed to settle to the surface of the earth; or to actually cover or roof in plants.

DEVICES FOR PREVENTING RAPID RADIATION OF HEAT.

Devices designed to prevent a rapid radiation of heat from the earth include screens which can be drawn over plants, vineyards, and groves; loose substances with which low plants may be covered; and smudge fires built to the windward of areas for which protection is

desired. These devices, which of a necessity are operative for very limited areas, have been described by Prof. W. H. Hammon and by Prof. A. G. McAdie.

Glass screens.—In greenhouses and hotbeds advantage is taken of the peculiar property of glass which allows the heat rays of the sun to pass through it, and is almost impervious to the dark *heat* rays from the earth and plants. This is one of the most perfect screens possible, since it not only prevents the loss of heat by radiation, but receives and retains the heat from the sun. The expense precludes its adoption except for the protection of valuable plants and flowers.

Screens of other solid materials have been quite extensivly used in protecting vineyards and citrus groves where intense cultivation is practiced and where the location of the groves near an excellent market admits of profit even with expensive methods of cultivation.

Cloth screens.—In Italy and portions of France screens made of muslin strung on wires stretched on poles above the tops of trees or vines have been used extensively. These screens are drawn on nights when frosts are probable and are pushed back during the day. When the season has advanced so far as to preclude further danger, they are taken down and stored. Of course such a plan could be operated only on a very limited scale, and would then be expensive. This plan has been recently successfully tried in the orange groves of southern California.

Lath screens.—During the past few years screens made of laths fastened to ordinary telephone wire (the spaces between them being about the width of the laths) have been extensively used in Florida. These are spread over a frame erected above the trees or plants. The screens serve not only as a fair protection from frost, but also as a shade from the hot sun. When no longer needed, they can be rolled up and stored away for preservation. At first thought it would seem improbable that a screen covering only half the space (the spaces being as wide as the laths) would afford much protection; but when it is considered that laths have considerable thickness it is plain that while only one-half the vertical rays are screened those inclined between the vertical and horizontal are partially intercepted by the edges as well as the faces of the laths. As a matter of fact, about three-fourths of the sky is screened by this means. By placing the laths in north and south directions the direct rays of the morning sun are completely cut off from the orchard, which admits of the temperature rising slowly. This reduces the liability of injury to plants.

Paper cover.—A new form of cover, known as an antifrost cover, has been devised in the San Francisco Weather Bureau office. It is, however, best suited for the protection of small fruit, garden flowers, and vegetables. It is an old and widely known practice to cover garden flowers with newspapers when frost is expected. In the antifrost cover there is used instead of a single cover a double layer of prepared paper with an intervening air space. This prevents almost perfectly the escape of the long heat waves from the ground. Such covering can be spread above the plant an hour or two before sunset, i. e., before the ground has lost much of its heat. If at the same time shallow pans of warm water are placed under the cover, an effective screen and serviceable supply of water are provided. It is also to be noted that we use this cover at a time when the roots are absorbing vigorously and transpiration by the leaves is at a maximum. We therefore prevent any lowering of temperature at the leaf surface and store up in the plant a quantity of heat energy for expenditure slowly through the night hours. The actual temperature of the air is of less importance than the temperature of the leaf or fruit surface. A deposit of moisture is advantageous.

A special use of the antifrost cover in connection with tree fruits is to unroll the covers in the orchard before sunset and to roll them up about 4 o'clock in the morning.

Other methods.—Strawberries and other low plants are frequently protected by covering them with straw or other loose substances.

Frequently young potato plants are saved by plowing a furrow alongside and allowing the dirt to bury them.

Cranberry growers in the marshes of Massachusetts, New Jersey, and Wisconsin flood the marshes when frost is expected. In this case the protection is probably due, for the most part, to the high specific heat of the water, as only portions of this land are submerged. Cultivating, draining, and sanding are now resorted to with good results.

DEVICES FOR ADDING HEAT TO THE AIR.

The first successful attempts to prevent injury by frost by the use of heating devices on a large scale occurred in California about 1896. Among a number of protective methods based upon heating, described by Prof. A. G. McAdie, may be mentioned the device of Edward Copely, of Riverside, Cal., in which a wire basket filled with coal and sufficient kindling to insure proper combustion was used. The method met with much favor in the orange groves of California. With from 20 to 40 of these baskets to the acre the temperature under ordinary winter weather conditions can be raised 3° or more.

The first use of oil that we have a record of was by Everett, at Arlington, Cal., and the first use of hot water was by Meacham at Riverside.

A severe frost about the end of December, 1895, caused great damage to the orange crop in the Riverside section, and the following year may be said to have marked the beginning of the frost-fighting campaign. The Riverside Horticultural Club took an active interest in the problem and many experiments were conducted by such men as those above mentioned and Koethen, Reed, Holmes, Hall, Hammon, McAdie, and others. The work has gone on, and there are now on the market many forms of heating devices. A convenient and serviceable frost alarm thermometer and oil pot were devised by the Weather Bureau official at Fresno, Cal., Mr. J. P. Bolton, for use in the vineyards during the spring frosts. There have recently been put upon the market in the apple sections of Colorado and other Rocky Mountain States orchard heaters burning oil, coal, or other fuel.

The various forms of heaters are known by the name of their inventors or are otherwise designated. It is not desirable in a bulletin of this character to advocate the use of any one special form. Indeed the heater that is most satisfactory in one locality and for a given crop may prove less satisfactory under changed conditions.

As showing the efficiency of these heaters, the following report upon a certain coal-burning heater was made by Prof. O. B. Whipple, field horticulturist, Colorado Experiment Station.

I have the following report to make on the coal-burning heater demonstration, held at your place on the evening of November 12, 1908:

The conditions under which the test was held were such as one would experience in fighting frost in the spring. One hundred heaters were well spaced at 29.5 feet, covering 2 acres of ground. In placing the thermometers, I was careful to see that no pots were closer than 20.5 feet. Waste saturated with one-half crude oil and one-half kerosene with tree prunings were used to start the fires. The fires were started at 7.45 and responded promptly, as the table will show. One man lighted the 100 pots in 8 minutes. All pots started without trouble. The pots were refilled, begining at 10.15, one man completing the job in 40 minutes. A few of the pots on a part of the block were again refilled at 12.30, and all burned until the end of the test. In all 2 tons of coal were consumed in the 7 hours, making a total of 40 pounds per pot. The coal used was one-half lump and one-half nut and came from the Farmers' mine and Stokes' mine. The temperature readings shown in the table were taken at an elevation of 7 feet from the ground. A thermometer at 12 feet from the ground inside the heated area showed 1° higher temperature.

The following table shows the temperature readings for the full 7 hours:

Hour.	Outside temperature.	Inside temperature.	Gain.	Hour.	Outside temperature.	Inside temperature.	Gain.
p. m.	°F.	°F.	°F.	*p. m.*	°F.	°F.	°F.
7.45	30.2	30.2	.....	12.00	29.3	36.9	7.6
8.15	31.1	35.6	4.5	*a. m.*			
8.30	28.4	32.9	4.5	12.30	25.7	32.6	6.9
9.00	28.4	34.2	5.8	1.00	26.1	32.4	6.3
9.30	28.4	33.8	5.4	1.30	26.6	33.8	7.2
10.00	28.4	34.2	5.8	2.00	28.4	35.6	7.2
10.30	27.5	32.9	5.4	2.30	28.4	34.5	6.1
11.00	27.5	33.8	6.3	3.00	28.4	33.4	5.0
11.30	28.4	36.0	7.6				

The following table is from the report of the committee in the "under actual frost conditions" test held at W. C. Knight's orchard, Grand Junction, Colo., October 28, 1908. The committee consisted of O. B. Whipple, field horticulturist Colorado Experiment Station; W. C. Knight and John M. Harvey, fruit growers:

Outside temperature.	Hour.	Inside temperature, with 50 coal heaters with concentrated draft.	Gain.	Inside temperature, with 100 oil heaters, best make.	Gain.	Inside temperature, with 80 coal heaters, open draft on all sides.	Gain.
°F.	*p. m.*	°F.	°F.	°F.	°F.	°F.	°F.
35	8.00	35	.....	35	.....	35	.....
35	8.15	38	3	38	3	36	.....
33	8.30	39	6	39	6	38	5
32	8.45	40	8	40	8	37	5
34	9.00	40	6	39	5	38	4
34	9.30	40	6	39	5	37	3
34	10.00	40	6	38	4	38	4
32	10.30	36	4	37	5	36½	4½
31	11.00	35	4	36	5	35	4
30	11.30	36½	6½	36	6	34	4
28	12.00	35½	7½	36	6	34	5
	a. m.						
27	12.30	34½	7½	35	8	32	5
27	1.00	34½	7½	35	8	32	5
Average gain.....			6	.....	5½	.....	4.4

The following account of the use of smudge pots, as they are called, is given by Mr. Richard H. Sullivan, local forecaster, in the Monthly Weather Review for 1910 (vol. 38, pp. 412–413):

The inclosed series of observations were taken for the benefit of members of the Sedgwick County Horticultural Society and others in connection with the frost warnings of March 30 and April 5 and tests of smudge pots in the 1-acre orchard of Mr. Albert Kunkel, in Wichita, on March 31 and April 6, and in a selected plat of the orchard of the Thomas Orchard Co., 3 miles west of Wichita, on April 6. The readings for the Kunkel orchard on March 31 and the Thomas orchard on April 6 were made by the writer, 2 minimum and 5 exposed thermometers, 1 anemometer, and 1 commercial thermometer being used.

On Mr. Kunkel's place 79 pots to the acre were used on March 31 and 70 pots to the acre on April 6; his fruit is still unharmed. In the Thomas orchard, 50 pots to the acre, or 500 pots in all, were used among Jonathan and Grimes Golden apple trees. Unfortunately, the temperature in the latter orchard could not be kept above freezing after 4 a. m. on account of lack of fuel oil. The small number of pots to the acre made it necessary to run them at nearly full capacity, and in order to carry the heat through to sunrise the lighting was delayed until 2 a. m. However, the manager does not consider his loss as great as in some orchards in the vicinity.

The figures show that with from 70 to 80 pots to the acre a fruit crop can be saved when the temperature falls to 25°, or even to 22°, if the work is done thoroughly and systematically. These are the first known practical tests of this character that have been made in this vicinity, and the whole proposition rests upon the question of how much expense for such insurance the investment can stand and still render a profit.

Where artificial methods of prevention of damage were not used during the freezes of March 31 and April 6, especially in the lowlands, there was general loss of apricots, peaches, plums, pears, and a very large proportion of apple buds, making the third series of disastrous spring frosts in four years. Last year the whole fruit-tree crop was killed by a single freeze during the night of April 30–May 1, warning of which was given the day before.

There is now a general disposition among local horticulturists to adopt the methods of commercial fruit growers elsewhere to prevent damage by frosts and freezes, and the belief is steadily growing that the warnings of this service must be heeded if loss is to be avoided.

Comparative temperature, wind velocity, and weather readings made in connection with test of central-draft smudge pots for prevention of damage by frost or freezing in the orchard of Albert Kunkel.

Half-hourly observations were taken from midnight of March 30–31 to 7.30 a. m. March 31. The arrangement and height of the various instruments are indicated by the letters in the diagram (fig. 3) and the explanation herewith: *A*, Weather Bureau anemometer, 5 feet above ground; *a*, W. B. minimum thermometer No. 6733, 3 feet 8 inches above ground; *b*, W. B. standard thermometer No. 4403, 4 feet 9 inches above ground; *c*, W. B. standard thermometer No. 4788, 4 feet 9 inches above ground; *d*, W. B. standard thermometer No. 4789, 4 feet 9 inches above ground; *e*, W. B. minimum thermometer No. 9436, 3 feet 8 inches above ground; *f*, W. B. standard thermometer No. 3135, 4 feet 9 inches above ground; *g*, W. B. standard thermometer No. 5671, 4 feet 9 inches above ground; *h*, commercial thermometer, 1 foot above ground; *i*, residence; *j*, greenhouse; *k*, barn.

The smudge pots, 79 in number, were lighted between 5.30 and 5.45 a. m. March 31, and extinguished at 7.30 a. m. The temperatures at 6 a. m. in the accompanying table show the effectiveness of such a method during frosty periods. Fuel oil, costing 4 cents per gallon, was used.



FIG. 3.—Diagram of Kunkel orchard, Wichita, Kans. Shaded areas show limits of smudge pots.

Winds—Northwesterly to 2.30 a. m.; northerly to 6.30 a. m.; northeasterly to 7.30 a. m.

TABLE 1.—*Temperature, weather, and wind velocity, Mar. 31, 1910.*

[Letters in heading indicate position of thermometers.]

Time.	*a*	*b*	*c*	*d*	*e*	*f*	*g*	*h*	Weather.	*A*
	°F.	°F.	°F.	°F.	°F.	°F.	°F.	°F.		Mile.
12.00 midnight	39.0	40.5	41.0	41.0	39.0	41.0	40.5	39.0	Clear	0.2
12.30 a. m.	38.5	40.0	39.5	39.5	38.5	39.5	39.5	38.0	do	.2
1.00 a. m.	36.5	38.5	38.0	38.0	37.0	38.5	38.5	37.0	do	.2
1.30 a. m.	35.5	37.5	37.0	37.0	36.0	38.0	37.5	36.0	do	.2
2.00 a. m.	35.0	37.0	37.0	36.0	35.5	36.0	36.0	34.5	do	.3
2.30 a. m.	34.0	36.0	35.5	35.5	34.0	35.5	35.5	34.0	do	.3
3.00 a. m.	35.5	37.0	37.5	37.0	36.0	37.5	37.0	35.0	do	.4
3.30 a. m.	36.0	37.0	39.0	39.0	37.0	38.5	38.0	36.0	do	.5
4.00 a. m.	36.5	37.0	36.5	36.0	35.5	37.0	37.0	36.0	do	.3
4.30 a. m.	35.5	36.5	36.0	36.0	35.0	36.5	36.0	35.0	do	.3
5.00 a. m.	33.0	34.0	33.5	33.0	33.0	34.0	33.0	31.0	do	.2
5.30 a. m.	[1]31.0	32.5	32.5	32.0	31.5	33.0	33.0	30.0	do	.2
6.00 a. m.	31.0	44.0	40.5	38.0	34.0	38.0	38.0	36.0	do	.3
6.30 a. m.	33.0	39.0	40.0	38.5	34.5	37.5	38.0	38.0	do	.4
7.00 a. m.	35.5	38.0	38.0	37.5	35.5	38.0	39.0	38.0	do	.5
7.30 a. m.	40.0	40.5	41.0	41.0	40.0	40.5	40.5	40.0	do	.5

[1] Lowest temperatures recorded, 30.5° between 5 and 5.30 a. m.

79 smudge pots lighted, beginning at 5.30 a. m. and ending at 5.45 a. m.

Second test made by Mr. Albert Kunkel. Number of pots lighted, 70.

TABLE 2.—*Temperature and weather table, Apr. 6, 1910.*

[Letters in heading indicate location of thermometers as shown on diagram of Mar. 31, 1910.]

Time.	*a*	*b*	*c*	Weather.
	°F.	°F.	°F.	
3.00 a. m.	30	32	.....	Clear.
3.30 a. m.	28	31	.....	Do.
4.00 a. m.	27	34	.....	Do.
4.30 a. m.	26	35	.....	Do.
5.00 a. m.	26	36	.....	Do.
5.30 a. m.	27	36	37	Do.
6.00 a. m.	28	38	38	Do.
6.30 a. m.	28	35	38	Do.
7.00 a. m.	30	38	38	Do.

70 smudge pots lighted, beginning at 3.30 a. m.

Comparative temperature, wind velocity, and weather readings made in connection with test of central-draft smudge pots for prevention of damage by frost or freezing in the orchard of the Thomas Orchard Co., 3 miles west of Wichita.

The smudged portion of the orchard was a selected 10-acre plat containing Jonathan and Grimes Golden apples; the pots were arranged 50 to the acre, or 500 in all. The instruments were placed at a height of 6 feet, except one commercial thermometer, which was located 1 foot above the ground near the center. The thermometers are arranged in the table in the order of reading. The pots were charged with 3 quarts of fuel oil, and lighting was delayed until 2 a. m. on account of lack of oil. The 500 pots were lighted between 2 and 2.35 a. m., and insufficient oil to hold the fires at reasonable capacity resulted in inability to keep temperature above freezing after 4 a. m., and by daylight practically all of the oil had been consumed.

Winds: Northwesterly to 4.30 a. m.; westerly after 4.30 a. m.

The letters in the diagram (fig. 4) of this plat indicate corresponding instruments in Table 3; numbers, United States Weather Bureau thermometers; *d*, commercial thermometer; *A*, anemometer.



FIG. 4.—Diagram of plat of Thomas Orchard Co., Wichita, Kans.

TABLE 3.—*Temperature, weather, and wind velocity.*

[Letters in heading indicate position of thermometers.]

Time.	*a*	*b*	*c*	*d*	*e*	*f*	*g*	*h*	Weather.	A
	°F.	°F.	°F.	°F.	°F.	°F.	°F.	°F.		Mile.
12.00 midnight	31.5	31.5	32.0	31.5	32.0	31.0	31.0	31.0	Clear	0.2
1.00 a. m.	31.0	31.0	31.0	32.0	31.5	31.0	31.0	30.5	do	.2
2.00 a. m.	30.0	[1]30.0	30.0	30.0	30.0	29.5	31.5	29.5	do	.1
3.00 a. m.	33.5	33.5	33.5	35.0	34.0	33.5	34.5	33.0	do	.1
4.00 a. m.	32.5	32.0	32.5	32.0	32.5	32.5	33.5	32.0	do	.1
5.00 a. m.	31.0	31.5	31.0	31.5	30.5	29.0	30.0	30.0	do	.2
6.00 a. m.	31.0	30.0	30.0	30.0	30.0	29.0	29.0	28.5	do	.1

[1] Lowest temperature recorded, 29.0° between 1 and 2 a. m.

500 smudge pots lighted, beginning at 2 a. m. and ending at 6.35 a. m.

Additional report, April 30, 1910.

Further trials in the Thomas orchard on April 19, 24–25, fully confirm the efficacy of the firing methods above explained in detail. The wind-velocity records show, as was to be expected, that the most damaging temperatures occur after the wind has died down. Artificial preventive methods can be used effectively despite the oft-repeated arguments that it is too windy in Kansas.

In a paper read before the California Fruit Growers' Convention at Watsonville, December, 1909, the district forecaster discusses in some detail the various forms of heating devices, such as open fires, smudge fires, coal baskets, oil pots, orchard heaters, etc., as used in the protection of citrus fruits in that State.

Many small fires are found to be much more effective than one or two large fires. The great mass of experiments made in California orchards show that direct heating of the air by open fires has not been sufficient to prevent injury at times of very low temperatures. A large amount of the heat thus produced is wasted; the efficiency of the method is low. This is illustrated by the following experience of a gentleman who is a close observer, an earnest student of the problem of frost protection, and one in whom I have the greatest confidence. During the night of December 20–21, 1908, on a certain California ranch, the temperature for 14 hours ranged between 19° and 24° F. For 36 hours the temperature did not rise above 28°. During this night the gentleman referred to burned 15 cords of wood and about 40 tons of wet hay in his efforts to protect his orchard. The relative humidity was low, there was little movement of the air, and he reports that "the smoke rose as straight as a pine tree." At the intersection of two roads in the orchard a large fire was maintained, and 30 feet distant the temperature in an olive tree was observed. Another fire was burning within 20 feet of this tree on a second side, and on yet a third side was a third fire maintained within 25 feet of the same tree. The temperature at the tree, however, remained at about 20° F. from 3 to 8 a. m. of December 21. This was the coldest weather in this locality since 1888. It is evident that in this case a large amount of heat escaped without producing the desired warming effect, i. e., was lost, wasted. It is, of course, well known that the rate of conduction of heat through air is low.

Antifrost candle.—A new form of heater, known as an antifrost candle, has been devised in the San Francisco Weather Bureau office. This consists of a tube of heavy cardboard or other suitable slow-burning material about 2 inches in diameter and 12 inches or more in length. The tube is filled with oil and cotton waste or wicking, or if sawdust is available this may be used. Suitable

stoppers and wicks close the ends of the tube. After filling, the candle is suspended by a wire about 3 feet below the fruit. It hangs in a horizontal direction or slightly tipped. A few inches above the candle a piece of sheet metal about 1 foot wide and 2 feet long is suspended. This metallic screen serves to prevent any possible burning by the open flame and catches the carbon particles, preventing their deposit on the fruit. The heat radiated from the upper surface of the metallic screen prevents a fall in temperature during the night and early morning hours. The great advantage of the candle is that the heat can be applied at the particular place where most needed and therefore a higher efficiency of the fuel used is obtained. Orchard heaters were a distinct advance upon the old-style open fires, but the new candles promise to be just as great an advance over the orchard heaters. In the new device less heat is wasted than in attempting to "warm up all out of doors."

DEVICES FOR ADDING MOISTURE TO THE AIR.

Smudge fires made of damp material have been more successfully used for the purpose of affording protection from frost than those made of tar, crude petroleum, and other dry materials, for the reason that in addition to creating a dense smoke they add to the atmosphere a considerable amount of evaporated water, which, though invisible, serves to retard the radiation of heat. When any considerable amount of water is thus converted into vapor and is distributed in the air which covers the area to be protected, a portion of it is likely to be condensed by the surrounding cooler air and appear in the form of mist, which acts as an agent to prevent the escape of heat from the earth; and the act of condensation sets free in the air some of the heat that has been expended in the process of evaporating the water contained in the smudge-fire materials. Thus the process is well ordered throughout. In the first place, an undue quantity of heat, whereby undesirable drafts would be created, is not communicated to the air by the fires, and the final result is that in reliberating, by condensation, the water in the air the particles of moisture remain suspended in the air, and some of the heat of the fires is returned to the air in a form which, to a certain degree, warms it without causing currents which would be calculated to disturb the smoke cloud, or pall, and the moist air which overlies the ground and upon whose presence the safety of the plants from frost depends.

Portable smudge fires.—A number of excellent devices have been tried in which the fires were built upon some vehicle by which they could be moved about the orchard. The advantages of this plan are several:

First. The fire can be moved to the section where most needed, which is generally along the windward side of the orchard.

Second. The loss of heat by an upward draft is almost entirely prevented, since the fire does not remain in one position long enough to establish such a draft. On this account much larger and consequently fewer fires, with equal efficiency, are possible.

Third. There is a much more uniform distribution of heat and smoke throughout the orchard.

Aside from the smudge fires, other devices have been adopted to charge the air directly with moisture, some of which are constructed on a plan too elaborate and expensive for general use.

The first operation involved is the evaporation of water in large quantities, either by means of evaporating pans or by the use of boilers with connecting pipes whereby steam can be generated, conducted, and discharged into the air in different parts of a field or orchard.

Spraying and sprinkling plants in times of threatened frost have also been attended with some degree of success. At the Everest ranch, at Riverside, Cal., sprinklers are placed at the top of 50-foot masts, which fill the air with a very fine spray. In nearly ever instance the protection against frost by this method has been sufficient.

Sprinkling gardens before sunrise on frosty mornings has proved of decided value.

Secondly, irrigation, which of necessity can be used only to a very limited extent and in certain favored localities, has been found to afford very complete protection from frost in fields, orchards, and vineyards which are equipped with irrigation ditches, and in the cranberry districts of Wisconsin and Cape Cod perfect protection from damage by frost is secured by flooding the cranberry marshes the day before heavy frost occurs in those sections.

In no case should water be allowed to stand in orchards, vineyards, or gardens during the early growing period while danger from frost is possible, for the reason that water unduly promotes the growth of young plants and thereby renders them more susceptible to damage by frost and cold than are the plants whose growth is retarded.

GENERAL OBSERVATIONS AND PRACTICAL RESULTS.

The general campaign of frost fighting, as thus far laid out, covers three main lines:

(1) Accurate advance information of the occurrence of frost.
(2) Applying preventive measures during critical hours.
(3) Preventing sudden warming of chilled fruit, or a too rapid thawing out.

Under the first of these heads great progress has been made in the past five years. Throughout the various fruit sections of the United States, Weather Bureau officials recognize the value of special study of types of frost maps. The work first undertaken in California on a large scale and for a distinct purpose is now an important feature of the forecaster's work at a large number of places. Frosts are found to occur as a consequence of certain movements of low and high pressure areas and the displacement of the lower air incident to

the pressure changes. Frost is a matter of air drainage, both on a large scale and on a small scale. In other words, frosts are successfully forecast because we understand the general movements of the lower air and in particular certain local circulations.

The forecaster anticipates a condition in which the lower air strata are quiet, dust free, and very dry. Under such condition there is rapid radiation and loss of heat from plant and ground and a sharp fall in temperature. While the forecaster gives the general conditions favorable for frost, it must be clearly understood that each individual orchardist must study his own problem of air movement in his own locality. These local circulations should be studied in connection with the daily weather map. It is also advisable to have certain instrumental records, such as are given by the thermograph and hygrograph.

Under the second head we have discussed the physical processes operative in the formation of frost and mentioned the various devices, such as smudging outfits, coal baskets, oil pots, orchard heaters, antifrost covers, candles, etc. Opinions will differ as to the relative efficiency of various devices, and some are undoubtedly more serviceable for certain localities than others.

There are three general principles used in frost-protection devices:

(1) Adding heat.
(2) Saving heat.
(3) Mixing or stirring the air.

Under the third head, methods based upon mixing or stirring the air, no special devices that are available and practicable are known. When nature mixes the air—i. e., on windy nights—frost does not occur. It is now known to meteorologists that layers of air of different temperature may lie close to one another without mixing. On nearly every frosty night there is a difference of 6°, 8°, or 10° in temperature between the ground and the air 10 feet above, the warmer layers being above. Where air is well mixed and there is good ventilation we seldom find frost.

WORK IN FLORIDA.

Devices used in Florida for the protection of fruit, vegetables, and berries from frost and light freezes conform in character to those tested in the California experiments, in which fires were found to be the most effective means of protection, and appliances for adding moisture to the air were successful only to a slight degree.

As artificial appliances are totally inadequate to add to the atmosphere any very appreciable amount of moisture, it is evident that methods which may have been found ineffective in the dry climate of California would possess value in localities where their office is confined to adding moisture to an already moist atmosphere. The

Gulf and South Atlantic Coast States, and, in fact, the country generally from the Mississippi Valley to the Atlantic seaboard, possess a moist atmosphere, and the fruits and tender vegetables of these districts can, therefore, be the more readily protected from frost by devices which add moisture to the air. As a means of adding moisture to the air, irrigation should be more effective than in California, and in localities where this method can be used protection will be assured except against hard freezes.

Owing to the comparatively inexpensive character of the materials used in damp smudge fires, they seem the best adapted for common use in groves and gardens. Berries and other low plants can be protected with but little expense by coverings of straw and other light materials. In Florida pine needles are successfully used to protect strawberries. Devices for actually heating the free and open air are expensive, and their value is dependent solely upon a comparatively still air and small, numerous, and well-distributed fires.

In Florida many experiments have been made with a view to adopting devices which will protect citrus trees and fruits from injury during the periods of severe cold which at times visit that section. The fact that these periods of cold are infrequent does not relieve the grower from the necessity of providing for their occurrence. Effective measures of protection necessarily vary with the conditions that confront the grower. Protection can be made effective by tents, coverings of cloth, sheds, and one-half and one-third lath shelters and their modifications, that retain heat from open wood fires and sheet-iron stoves. The expense of these methods is obviously too great to make their employment, except for choice varieties of fruit, commercially profitable.

In this State protection of citrus trees and fruit in the open field is usually afforded by what is known as "firing." This method is effective, except in the presence of severe freezes, provided rain or strong winds do not accompany the cold. Heavy rain will, of course, extinguish the fires, and strong winds will not permit a necessary accumulation of warm air about the trees. When no rain is falling and the wind is light, fires must be built on the windward side of the trees, so that the heat will be carried through the trees, and the number of the fires must be regulated by the degree of cold. In this form of protection the grower must lay in and place his materials for "firing" before the winter season sets in, and must, during the season, be on guard to utilize the materials (usually wood) that he has accumulated. When local indications point to dangerously low temperatures, or when warnings have been received from the Weather Bureau, he should watch his thermometers, exposed in the groves, and be prepared to "fire" with a rush when the mercury sinks to the danger point, 28° F., with a downward tendency. There is no

time on the eve of a freeze to collect firing materials, and the grower must not desert his grove for bed or board so long as danger by cold continues. He will have his hands full keeping his fires going, even if he has sufficient material for that purpose. Material should be available for at least three nights. Cold spells seldom last longer. The danger increases with the length of the cold spell. Trees and fruit become chilled, and the last night of a cold spell is likely to be the one that does the damage. The freeze of December 23–27, 1906, lasted five nights, and in some sections the last night produced the lowest temperatures of the period. In some localities growers had exhausted their supply of firing material and, as a consequence, suffered a loss of fruit and young trees.

In cases of severe cold, fruit can not be saved, and trees can be preserved only by banking up the trunks by sand or soil. Small trees can be buried, and older trees can at least be protected above the parts where they have been budded.

The damage that will result from a given degree of cold varies with the condition and size of the tree. Well-grown trees in a dormant condition will stand without serious injury, except to foliage, a temperature of 18° to 20°, provided it is not too long continued. When the sap is flowing these temperatures are likely to kill growing trees, and are apt to cut small, unprotected trees to the ground. Well-grown trees in a dormant condition will stand without injury a temperature of 26° of continued cold, and can for several hours stand temperatures of 22° to 24°. When the sap is flowing these temperatures continued for several hours will be more or less injurious. Cold that will freeze fruit will injure small trees. In the freeze of December 23–27, 1906, temperatures in parts of the orange belt fell to 20°, and even to 18° and below, on one and in some cases two nights without injury to large trees, except to foliage, and with a surprisingly small loss of fruit and small trees. This freeze had been preceded by about two months of rainless weather, and found the trees in a dormant condition. The destructive freezes of preceding years usually followed periods of wet weather that promoted a flow of sap.

WORK IN MISSOURI.

In the Monthly Weather Review for May, 1910, Mr. George Reeder, section director, expresses the following views relative to critical temperatures and the necessity of definite and detailed advance information in connection with frosts:

The horticulturists at the University of Missouri have closely determined the temperatures representing the danger point to fruit for the different stages in its development, as shown by the following brief summary of Circular No. 13, Missouri Experiment Station, 1909:

"Fully dormant peach buds can stand 8° or 9° below zero (F.). When they are appreciably swollen, zero is the danger point. When the buds are showing

pink, they can stand 15° above zero. When the buds are almost open, 25° is the point of danger. When the petals are beginning to fall, 28° above zero is cold enough to cause uneasiness. When the petals are off, they can stand 30° above zero. When the 'shucks' (calyx tubes) are beginning to fall off, 32° above zero is the danger point."

It is estimated that the April, 1910, freeze caused a loss to the people of Missouri of $2,500,000, and it would mean thousands of dollars to the fruit growers in carrying out protective measures in the future to know just when to start the heaters in the orchards. It has been demonstrated that it is not a difficult matter, if one is prepared for it, and knows how to start the work, to raise the temperature in an orchard 5° to 6°, sufficient during almost any cold night in the spring to save many an orchard from injury. The question asked so often is, "When shall we start the heaters?" To start them too soon is expensive, but to start them too late would be more expensive in the end. In this matter of advice the Weather Bureau must shoulder the burden, and it must take up the task with an earnestness that should prove successful. The forecaster must be thoroughly familiar, not only with the geographical location and topographical conditions of the section for which forecasts are made, but he must keep himself fully informed as to the progress of the fruit crop in its different stages of advancement. The State forecast issued by the district forecaster is all that is necessary for the public in general, but where specific information is desired it is not satisfactory. It will not suffice to simply send a weather message to a fruit locality in time of danger from low temperature, reading "Fair to-night and colder, with frost"; but the message must say how cold it is expected to be; for instance, something like the following: "Fair and colder to-night, with frost; temperature about 36°; no danger to fruit." And again, "Fair and colder to-night, with freezing temperature; 28° or lower by morning; start heaters after midnight."

While specifying the exact degree of cold expected may not be practicable for the district forecaster, owing to the extent of territory forecast for, and the limited time in which he has to make the forecast, it is being done, and can doubtless be brought to a high degree of accuracy, by the local forecasters.

DISTRIBUTION OF FORECASTS IN CALIFORNIA.

The system of frost warnings has reached a high degree of efficiency in California. The district forecaster sends telegrams each night to important fruit centers during the frost period, indicating the type of frost, whether light, heavy, or killing, the general weather conditions, and advice as to whether fire will be necessary.

To illustrate the way in which the information is distributed from the larger centers, the following statement is furnished by Mr. A. B. Wollaber, local forecaster at Los Angeles:

In California, where the changes in weather conditions can be only forecast for a short time in advance and where changes from one extreme to the opposite are often sudden, it has been found necessary to adopt a system whereby the local office can keep in touch at all times with the citrus industry. Here we find that the best means for accomplishing this purpose is the telephone, and in addition to the free distribution of our warnings by the long-distance companies operating in this section we have an arrangement whereby we can reach the entire fruit belt within half an hour and disseminate our warnings through the medium of the fruit exchanges. The Weather Bureau pays for this service and therefore is in a position to demand and to receive prompt

service. To accomplish this we have prepared a list of the exchanges in the fruit belt to which we send our warnings, and this list is on file with the chief operators of the long-distance companies. When we desire to send a warning we call for long-distance chief operator and give the message, which is immediately sent out. This system is used only in extreme cases, as when the regular maps and cards have been sent without a frost warning. That could occur in this manner: When a storm which gives indication of clouds or rain during the period covered by the morning forecast passes eastward more rapidly than was anticipated, causing a rapid clearing, and when later information from the district forecaster shows the probability of more severe conditions than anticipated on the morning reports.

Now, it happens in this section that not all ranches need protection at all times. The managers of the several fruit exchanges are personally familiar with conditions in their respective sections and are considered to be best qualified to disseminate this important information.

Special frost-warning cards are mailed to all post offices, fruit exchanges, and railroad depots before the frost period begins. There has grown up a personal relationship between the growers and the local office, and through this much good has been accomplished.

ORCHARD HEATING IN INDIANA.

In Monthly Weather Review, January, 1911, Mr. William M. Walton, jr., gives an interesting account of an effort to protect orchards under ordinary weather conditions in Indiana. Heaters holding 5 quarts of oil were placed in the orchard on March 31, 21 feet apart each way, with a small amount of straw on top. The first frost occurred April 6 at 6.30 p. m. Three persons lit 500 fires in about 30 minutes and within half an hour the temperature was 40° F., and alternate heaters were extinguished. In the unheated section of the orchard the temperature fell to 27° F., but in no place in the heated area did it get below 32°. On the 7th similar operations were carried on, but the heaters were kept going until 6.30 a. m. As the sun came up everything outside the heated area was white with frost and ice one-fourth of an inch in thickness had formed, while no frost whatever could be found in the orchard. Mr. Walton further states that on April 12 a heavy frost appeared at 11 p. m., but it was only necessary to burn alternate heaters.

Snow fell throughout the night of April 17, but as the temperature did not fall below freezing the fires were quenched a short time after they were started. Not a single blossom had been injured within the heated area previous to the 22d. Currants hung in full clusters with 10 or 11 on each stem. In the sections where heat had not been used blossoms were damaged to a considerable extent, the effect being most noticeable on currants, where it was impossible to find more than 5 or 6 on a stem.

At this point it is proper to refer to the weather conditions in Indiana during March and April. The mean temperature for March was the highest on record and vegetation was fully two weeks ahead

of the normal. On the other hand, during the latter half of April all previous records of low temperatures for that time of the year were broken and unusual conditions resulted.

In the words of Mr. Walton:

On April 22 a high northwest wind was blowing with temperature at 32°. It was imposible to get the heat up into the trees with the wind blowing a gale. However, we held our own quite well until daylight, when the temperature started downward. By 10 o'clock the thermometer read 25° and the wind was still blowing hard. We kept the heaters going, but the blossoms soon became like wax. We maintained the fight until the arrival of the weather forecast, which was for continued high winds and freezing temperature. At 11 a. m. nearly everything within the heated area as well as without was frozen and caused us to forego our efforts after having burned 3,300 gallons of oil. The freezing weather continued throughout the day and night of the 23d, 24th, and 25th, and was followed by heavy snow on the 26th. As a result we picked no gooseberries or currants from the 18 acres which were in bearing, nor any cherries, plums, or peaches, and but a few apples from the remainder of the orchard.

Having lost all tree and bush fruits, we turned our attention to 1⅝ acres of strawberries that were not in blossom during the period of severe weather just described. We placed 180 heaters in this plat and went through three nights of frost with temperature down to 25° and lower. On the night of May 13 we started the fires at 8.30 p. m. and kept them going until 7 a. m. It was difficult to prevent the frost from settling on the vines, as there was no foliage above to assist in retaining the heat, as in the orchard. However, we went through the strawberry season without rain, and yet harvested a larger crop of berries than we did the previous year from a larger tract under more favorable moisture conditions. This goes to show that the heaters should at least share in the credit for the good results obtained.

We burned during the season about 4,000 gallons of fuel oil, which would have been sufficient in any ordinary year, or, at least, would have saved the crop had it been used during the spring of 1908 and 1909 instead of 1910.

We went into the matter this year to fight the spring frosts, but when winter entered the fray we were compelled to surrender. We are not discouraged by the failure, however, and expect to try again. The experience gained is particularly valuable, and gives us increased energy for further experiment. It also convinces us that with proper equipment we can save our entire crop during seasons having ordinary weather conditions in the spring.

WORK IN OREGON.

In Farmers' Bulletin No. 401, Mr. P. J. O'Gara describes in detail the protection of orchards in the Pacific Northwest from spring frosts by means of fire and smudges. The same author, in the Monthly Weather Review for September, 1910, describes work in the Rogue River Valley during the spring of 1910. The work has been extensively carried on, and the following brief quotations will serve to show what has been done:

During the past season there were just a few orchards in the danger zone carelessly left without protection; luckily these have served as a check on our work. In every case the unprotected orchards lying within the frost zone were

badly damaged and in some cases the crop entirely destroyed, while the protected orchards did not show a frost mark or ring on any of the fruit. * * *

It is believed by the writer, as well as the growers, that the Rogue River Valley has not only settled the problem of orchard heating, but also the important matter of accurately forecasting the frost. During the past two years not a single serious error has been made in forecasting. Whether this can be done indefinitely remains to be seen, but the writer believes that it is entirely possible. The writer does not attempt to say that the same methods will apply in other localities, having entirely different conditions, but it is believed that for every locality the optimum time for making the forecasts may be found, as well as all the other factors which enter into the problem. This at least is worth a trial.

WORK IN UTAH.

Variability of frost injury on fruit buds.—Dr. P. F. Homer in the Monthly Weather Review for April, 1911, gives an extremely interesting review of observations made in Utah orchards to determine, if possible, the factors that control the resistance of fruit buds to temperatures below freezing. The author states:

The injury done by frost takes place in two ways. When the cold is not too intense the ice forms in the intercellular spaces, and under the influence of a sudden rise of temperature the moisture evaporates directly into the atmosphere and the cells dry out and shrivel up for lack of moisture. In the other case, under the influence of severe cold the cell sap is frozen within the cell, the protoplasmic membranes are ruptured, and often the cell walls, by the expansion of the freezing liquid. The degrees of harm that any given temperature may do probably depend on the concentration of the cell sap, and the factors which influence this are probably the controlling ones in determining injury.

The author shows that fogs, smudges, cloudiness in the mornings, and any condition that prevents a rapid thawing of the frozen tissue lessen the injury; while conditions that accelerate the thawing and drying increase the injury. Such injurious conditions are found at Pleasant Grove in the cold, drying north winds that follow the April freezes.

Orchards situated on the western slope will escape injury when those on the eastern slope will lose all their fruit. It is almost impossible to grow peaches on the west side of Utah Valley, while orchards close up against the western slope of the Wasatch Mountains, only 12 miles distant, are among the finest in the State.

The author gives a detailed account of the spring weather for 1907–1910, showing how the peach and apricot yields were affected.

Dr. Homer concludes:

There is a great variation in the amount of cold a bud or blossom can stand, but there is also great variation in the individual buds in their power to withstand cold. Thus a freeze that will kill one bud on a twig will leave unharmed the next three and then perhaps kill another, all apparently in the same state of advancement. One bud may be spared in a cluster of apples or pears or

cherries and all the rest taken. So it is sometimes with trees. One tree may bear a full crop and one in another row be fruitless. In many cases on Provo Bench an orchard on one side of the highway will have a good crop of fruit while one on the other side of the road will not have one live bud in several hundred. This seems to be true in orchards heated and not heated, pruned and unpruned, in sod or cultivated. Several unheated orchards will have more fruit than some that were heated.

There is much work to be done in determining the resistance of the fruit buds to frost and the factors that bring about such remarkable difference.

MISCELLANEOUS.

A TWO YEARS' STUDY OF SPRING FROSTS AT WILLIAMSTOWN, MASS.

A valuable paper by Prof. W. I. Milham on the above subject was published in the Monthly Weather Review for August, 1908. The special purpose of the investigation was the critical study of the spring frosts, first, to test the prevalent belief that the probable minimum temperature could be computed from the maximum and dew point, or from the reading of the wet-bulb thermometer of the afternoon before the frost; second, to study the variation in the severity of frost at different places within a small area; and, third, to study the variation in the severity of frost with varying distances above the ground. Prof. Milham came to the following conclusions:

(1) The cool nights of spring when a frost might be expected are very dry, and the dew point lies so low that it plays practically no part in determining the minimum temperature.

(2) The amount of the drop from the maximum to the following minimum is very far from a constant, even if the characteristics of these nights seem very nearly the same. In each case in estimating the probable drop one must take into account the probable amount of cloud, the probable wind velocity, the possibility of a change in wind direction, and possibly other things.

Milham holds that two processes operate to produce the cooling, first, the importation of cooler air, and, second, the radiation of heat from the ground and the cooling of the air next to it by conduction. It might seem that the cool nights during which frosts seem probable could be divided into two groups, depending upon whether the importation of cold air or radiation from the ground was playing the larger part in producing the cooling. It is found, however, that both processes are nearly always operative. It may not be out of place at this point to call attention to the fact that the heat which has been absorbed by the ground is re-emitted as radiation of long wave lengths, from 4 μ to 20 μ.

Milham calls attention to the fact that the temperature of vegetation in the open, near the ground, in the coldest part of the tract may be 11° lower than the estimated minimum in a given shelter.

An interesting paper upon temperature variations near the surface of the soil during the formation of frost, by Mr. Dewey A. Seeley, local forecaster, can be found in the Monthly Weather Review for August, 1908. He is of the opinion that a considerable amount of the heat given off, during the condensation of vapor as frost and dew, is retained in saturated air near the ground level and also by the foliage of the plants themselves. The influence of a surface covering of vegetation on the temperature of the soil was marked. It was found that the soil beneath the grass was warmer both at night and during the day than the naked soil; but the temperature of the air above the grass was lower in both cases than over bare ground.

The dew-point as recorded at 7 p. m. and the minimum temperature registered the following morning for 60 days were tabulated and the differences computed. From these it appears that on the average the minimum temperature went 2.9° below the 7 p. m. dew point. In but three cases did the temperature fall more than 10° below the dew point recorded at 7 p. m. The thermograph traces made during clear weather with light wind show that the temperature fall goes on much more rapidly during the first part of the night, before the dew point is reached.

[A list giving the titles of all Farmers' Bulletins available for distribution will be sent free upon application to a Member of Congress or the Secretary of Agriculture.]

XUM

FARMERS' BULLETINS.

These bulletins are sent free of charge to any address upon application to the Secretary of Agriculture, Washington, D. C. Only the following are available for distribution:

No. 16. Leguminous Plants for Green Manuring and for Feeding. No. 19. Important Insecticides: Directions for their Preparation and Use. No. 21. Barnyard Manure. No. 22. Feeding Farm Animals. No. 23. Foods: Nutritive Value and Cost. No. 24. Hog Cholera and Swine Plague. No. 25. Peanuts: Culture and Uses. No. 26. Sweet Potatoes: Culture and Uses. No. 27. Flax for Seed and Fiber. No. 28. Weeds; and How to Kill Them. No. 29. Souring of Milk and Other Changes in Milk Products. No. 30. Grape Diseases on the Pacific Coast. No. 31. Alfalfa, or Lucern. No. 32. Silos and Silage. No. 33. Peach Growing for Market. No. 34. Meats: Composition and Cooking. No. 35. Potato Culture. No. 36. Cotton Seed and Its Products. No. 37. Kafir Corn: Characteristics, Culture, and Uses. No. 38. Spraying for Fruit Diseases. No. 39. Onion Culture. No. 40. Farm Drainage. No. 41. Fowls: Care and Feeding. No. 42. Facts About Milk. No. 43. Sewage Disposal on the Farm. No. 44. Commercial Fertilizers. No. 45. Some Insects Injurious to Stored Grain. No. 46. Irrigation in Humid Climates. No. 47. Insects Affecting the Cotton Plant. No. 48. The Manuring of Cotton. No. 49. Sheep Feeding. No. 50. Sorghum as a Forage Crop. No. 51. Standard Varieties of Chickens. No. 52. The Sugar Beet. No. 53. How to Grow Mushrooms. No. 54. Some Common Birds in Their Relation to Agriculture. No. 55. The Dairy Herd: Its Formation and Management. No. 56. Experiment Station Work—I. No. 57. Butter Making on the Farm. No. 58. The Soy Bean as a Forage Crop. No. 59. Bee Keeping. No. 60. Methods of Curing Tobacco. No. 61. Asparagus Culture. No. 62. Marketing Farm Produce. No. 63. Care of Milk on the Farm. No. 64. Ducks and Geese. No. 65. Experiment Station Work—II. No. 66. Meadows and Pastures. No. 67. Forestry for Farmers. No. 68. The Black Rot of the Cabbage. No. 69. Experiment Station Work—III. No. 70. The Principal Insect Enemies of the Grape. No. 71. Some Essentials of Beef Production. No. 72. Cattle Ranges of the Southwest. No. 73. Experiment Station Work—IV. No. 74. Milk as Food. No. 75. The Grain Smuts. No. 76. Tomato Growing. No. 77. The Liming of Soils. No. 78. Experiment Station Work—V. No. 79. Experiment Station Work—VI. No. 80. The Peach Twig-borer—an Important Enemy of Stone Fruits. No. 81. Corn Culture in the South. No. 82. The Culture of Tobacco. No. 83. Tobacco Soils. No. 84. Experiment Station Work—VII. No. 85. Fish as Food. No. 86. Thirty Poisonous Plants. No. 87. Experiment Station Work—VIII. No. 88. Alkali Lands. No. 89. Cowpeas. No. 90. The Manufacture of Sorghum Sirup. No. 91. Potato Diseases and Their Treatment. No. 92. Experiment Station Work—IX. No. 93. Sugar as Food. No. 94. The Vegetable Garden. No. 95. Good Roads for Farmers. No. 96. Raising Sheep for Mutton. No. 97. Experiment Station Work—X. No. 98. Suggestions to Southern Farmers. No. 99. Common Insects on Shade Trees. No. 100. Hog Raising in the South. No. 101. Millets. No. 102. Southern Forage Plants. No. 103. Experiment Station Work—XI. (In press.)